



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07K 14/47, A01K 67/027, A61K 38/08, C07K 7/06, C07K 16/40, C12N 1/19, C12N 1/21, C12N 5/16, C12N 9/64, C12N 15/57, C12Q 1/68, G01N 33/50	A2	(11) International Publication Number: WO 00/47618 (43) International Publication Date: 17 August 2000 (17.08.2000)
(21) International Application Number: PCT/US00/03819 (22) International Filing Date: 10 February 2000 (10.02.2000) (30) Priority Data: 60/119,571 10 February 1999 (10.02.1999) US 60/139,172 15 June 1999 (15.06.1999) US (60) Parent Application or Grant ELAN PHARMACEUTICALS, INC. [/]; (). ANDERSON, John, P. [/]; (). BASI, Guriqbal [/]; (). DOANE, Minh, Tam [/]; (). FRIGON, Normand [/]; (). JOHN, Varghese [/]; (). POWER, Michael [/]; (). SINHA, Sukanto [/]; (). TATSUNO, Gwen [/]; (). TUNG, Jay [/]; (). WANG, Shuwen [/]; (). MCCONLOGUE, Lisa [/]; (). ANDERSON, John, P. [/]; (). BASI, Guriqbal [/]; (). DOANE, Minh, Tam [/]; (). FRIGON, Normand [/]; (). JOHN, Varghese [/]; (). POWER, Michael [/]; (). SINHA, Sukanto [/]; (). TATSUNO, Gwen [/]; (). TUNG, Jay [/]; (). WANG, Shuwen [/]; (). MCCONLOGUE, Lisa [/]; (). STRATFORD, Carol, A. ; ().		Published
(54) Title: 'beta'-SECRETASE ENZYME COMPOSITIONS AND METHODS (54) Titre: COMPOSITIONS A BASE D'ENZYME BETA-SECRETASE ET PROCEDES (57) Abstract <p>Disclosed are various forms of an active, isolated 'beta'-secretase enzyme in purified and recombinant form. This enzyme is implicated in the production of amyloid plaque components which accumulate in the brains of individuals afflicted with Alzheimer's disease. Recombinant cells that produce this enzyme either alone or in combination with some of its natural substrates ('beta'-APPwt and 'beta'-APPsw) are also disclosed, as are antibodies directed to such proteins. These compositions are useful for use in methods of selecting compounds that modulate 'beta'-secretase. Inhibitors of 'beta'-secretase are implicated as therapeutics in the treatment of neurodegenerative diseases, such as Alzheimer's disease.</p> (57) Abrégé <p>L'invention concerne différents types d'enzyme bêta-sécrétase active isolée, sous forme purifiée et de recombinaison, intervenant dans la production des éléments de plaque amyloïde qui s'accumulent dans le cerveau des personnes souffrant de la maladie d'Alzheimer. L'invention concerne en outre des cellules de recombinaison qui produisent l'enzyme en question, séparément ou en combinaison avec certains des ses substrats naturels ('beta'-APPwt et 'beta'-APPsw), et elle concerne aussi des anticorps vis-à-vis de ces protéines. Les compositions considérées sont utiles dans les procédés relatifs à la sélection des composés qui modulent la bêta-sécrétase. Les inhibiteurs de la bêta-sécrétase interviennent comme éléments thérapeutiques dans le traitement des maladies neurodégénératives comme la maladie d'Alzheimer.</p>		

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/47, C12N 9/64, C07K 7/06, 16/40, C12N 15/57, 5/16, 1/21, 1/19, G01N 33/50, A01K 67/027, A61K 38/08, C12Q 1/68		A2	(11) International Publication Number: WO 00/47618 (43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/US00/03819 (22) International Filing Date: 10 February 2000 (10.02.00) (30) Priority Data: 60/119,571 10 February 1999 (10.02.99) US 60/139,172 15 June 1999 (15.06.99) US (71) Applicant (for all designated States except US): ELAN PHARMACEUTICALS, INC. [-/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ANDERSON, John, P. [-/US]; 21 Bucareli Drive, South San Francisco, CA 94132 (US). BASI, Guriqbal [-/US]; 514 Rhoades Drive, Palo Alto, CA 94303 (US). DOANE, Minh, Tam [-/US]; 24003 Malibu Road, Hayward, CA 94545 (US). FRIGON, Normand [-/US]; 201-C Richmond Drive, Milbrae, CA 94030 (US). JOHN, Varghese [-/US]; 1772 18th Avenue, San Francisco, CA 94122 (US). POWER, Michael [-/US]; 4263 Blue Ridge Street, Fremont, CA 94536 (US). SINHA, Sukanto [-/US]; 808 Junipero Serra Drive, San Francisco, CA 94127 (US). TATSUNO, Gwen [-/US]; 5910 Pinewood		Road, Oakland, CA 94611 (US). TUNG, Jay [-/US]; 2224 Semeria Avenue, Belmont, CA 94002 (US). WANG, Shuwen [-/US]; 1226 12th Avenue #2, San Francisco, CA 94122 (US). MCCONLOGUE, Lisa [-/US]; 283 Juanita Way, San Francisco, CA 94127 (US). (74) Agent: STRATFORD, Carol, A.; Elan Pharmaceuticals, Inc., 800 Gateway Boulevard, South San Francisco, CA 94080 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: β -SECRETASE ENZYME COMPOSITIONS AND METHODS



(57) Abstract

Disclosed are various forms of an active, isolated β -secretase enzyme in purified and recombinant form. This enzyme is implicated in the production of amyloid plaque components which accumulate in the brains of individuals afflicted with Alzheimer's disease. Recombinant cells that produce this enzyme either alone or in combination with some of its natural substrates (β -APPwt and β -APPsw) are also disclosed, as are antibodies directed to such proteins. These compositions are useful for use in methods of selecting compounds that modulate β -secretase. Inhibitors of β -secretase are implicated as therapeutics in the treatment of neurodegenerative diseases, such as Alzheimer's disease.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Lichtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

5

β-SECRETASE ENZYME COMPOSITIONS AND METHODS

10

5 Field of the Invention

15

20

20

25

15

The invention relates to the discovery of various active forms of β-secretase, an enzyme that cleaves β-amyloid precursor protein (APP) at one of the two cleavage sites necessary to produce β-amyloid peptide (Aβ). The invention also relates to inhibitors of this enzyme, which are considered candidates for therapeutics in the treatment of amyloidogenic diseases such as Alzheimer's disease. Further aspects of the present invention include screening methods, assays, and kits for discovering such therapeutic inhibitors, as well as diagnostic methods for determining whether an individual carries a mutant form of the enzyme.

Background of the Invention

30

20

35

40

25

45

Alzheimer's disease is characterized by the presence of numerous amyloid plaques and neurofibrillary tangles present in the brain, particularly in those regions of the brain involved in memory and cognition. β-amyloid peptide (Aβ) is a 39-43 amino acid peptide that is major component of amyloid plaques and is produced by cleavage of a large protein known as the amyloid precursor protein (APP) at a specific site(s) within the N-terminal region of the protein. Normal processing of APP involves cleavage of the protein at point 16-17 amino acids C-terminal to the N-terminus of the β-AP region, releasing a secreted ectodomain, α-sAPP, thus precluding production of β-AP. Cleavage by β-secretase enzyme of APP between Met⁶⁷¹ and Asp⁶⁷² and subsequent processing at the C-terminal end of APP produces Aβ peptide, which is highly implicated in the etiology of Alzheimer's pathology (Seubert, *et al.*, in *Pharmacological Treatment of Alzheimer's disease*, Wiley-Liss, Inc., pp. 345-366, 1997; Zhao, J., *et al.* J. Biol. Chem. 271: 31407-31411, 1996).

50

30

It is not clear whether β-secretase enzyme levels and/or activity is inherently higher than normal in Alzheimer's patients; however, it is clear that its cleavage product, Aβ peptide, is abnormally concentrated in amyloid plaques present in their brains. Therefore, it would be desirable to isolate, purify and characterize the enzyme responsible for the

55

5 pathogenic cleavage of APP in order to help answer this and other questions surrounding the
etiology of the disease. In particular, it is also desirable to utilize the isolated enzyme, or
active fragments thereof, in methods for screening candidate drugs for ability to inhibit the
10 activity of β -secretase. Drugs exhibiting inhibitory effects on β -secretase activity are
5 expected to be useful therapeutics in the treatment of Alzheimer's disease and other
amyloidogenic disorders characterized by deposition of A β peptide containing fibrils.

15 U.S. Patent 5,744,346 (Chrysler, *et al.*) describes the initial isolation and partial
purification of β -secretase enzyme characterized by its size (apparent molecular weight in the
range of 260 to 300 kilodaltons when measured by gel exclusion chromatography) and
20 enzymatic activity (ability to cleave the 695-amino acid isotype of β -amyloid precursor
protein between amino acids 596 and 597). The present invention provides a significant
improvement in the purity of β -secretase enzyme, by providing a purified β -secretase enzyme
that is at least 200 fold purer than that previously described. Such a purified protein has utility
25 in a number of applications, including crystallization for structure determination. The
15 invention also provides methods for producing recombinant forms of β -secretase enzymes
that have the same size and enzymatic profiles as the naturally occurring forms. It is a further
discovery of the present invention that human β -secretase is a so-called "aspartyl" (or
30 "aspartic") protease.

20 Summary of the Invention

35 This invention is directed to a β -secretase protein that has now been purified to
apparent homogeneity, and in particular to a purified protein characterized by a specific
activity of at least about 0.2×10^5 and preferably at least 1.0×10^5 nM/h/ μ g protein in a
40 representative β -secretase assay, the MBP-C125sw substrate assay. The resulting enzyme,
25 which has a characteristic activity in cleaving the 695-amino acid isotype of β -amyloid
precursor protein (β -APP) between amino acids 596 and 597 thereof, is at least 10,000-fold,
preferably at least 20,000-fold and, more preferably in excess of 200,000-fold higher specific
45 activity than an activity exhibited by a solubilized but unenriched membrane fraction from
human 293 cells, such as have been earlier characterized.

30 In one embodiment, the purified enzyme is fewer than 450 amino acids in length,
50 comprising a polypeptide having the amino acid sequence SEQ ID NO: 70 [63-452]. In
preferred embodiments, the purified protein exists in a variety of "truncated forms" relative

5 to the proenzyme referred to herein as SEQ ID NO: 2 [1-501], such as forms having amino acid sequences SEQ ID NO: 70 [63-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 58 [46-452], SEQ ID NO: 74 [22-452], SEQ ID
10 NO: 58 [46-452]. More generally, it has been found that particularly useful forms of the enzyme, particularly with regard to the crystallization studies described herein, are characterized by an N-terminus at position 46 with respect to SEQ ID NO: 2 and a C-terminus between positions 452 and 470 with respect to SEQ ID NO: 2. and more
15 particularly, by an N-terminus at position 22 with respect to SEQ ID NO: 2 and a C-terminus between positions 452 and 470 with respect to SEQ ID NO: 2. These forms are considered to be cleaved in the transmembrane "anchor" domain. Other particularly useful
20 purified forms of the enzyme include: SEQ ID NO: 43 [46-501], SEQ ID NO: 66 [22-501], and SEQ ID NO: 2 [1-501]. More generally, it is appreciated that useful forms of the enzyme have an N-terminal residue corresponding to a residue selected from the group consisting of residues 22, 46, 58 and 63 with respect to SEQ ID NO: 2 and a C-terminus selected from a
25 residue between positions 452 and 501 with respect to SEQ ID NO: 2 or a C-terminus between residue positions 452 and 470 with respect to SEQ ID NO: 2. Also described herein are forms of enzyme isolated from a mouse, exemplified by SEQ ID NO: 65.

30 This invention is further directed to a crystalline protein composition formed from a purified β -secretase protein, such as the various protein compositions described above. According to one embodiment, the purified protein is characterized by an ability to bind to the β -secretase inhibitor substrate P10-P4'sta D \rightarrow V which is at least equal to an ability
35 exhibited by a protein having the amino acid sequence SEQ ID NO: 70 [46-419], when the proteins are tested for binding to said substrate under the same conditions. According to another embodiment, the purified protein forming the crystallization composition is characterized by a binding affinity for the β -secretase inhibitor substrate SEQ ID NO: 72 (P10-P4'sta D \rightarrow V) which is at least 1/100 of an affinity exhibited by a protein having the amino acid sequence SEQ ID NO: 43 [46-501], when said proteins are tested for binding to said substrate under the same conditions. Proteins forming the crystalline composition may be glycosylated or deglycosylated.

45 30 The invention also includes a crystalline protein composition containing a β -secretase substrate or inhibitor molecule, examples of which are provided herein, particularly exemplified by peptide-derived inhibitors such as SEQ ID NO: 78, SEQ ID NO: 72, SEQ ID
50

5 NO: 81, and derivatives thereof. Generally useful inhibitors in this regard will have a K_i of no more than about 50 μ M to 0.5 mM.

10 Another aspect of the invention is directed to an isolated protein, comprising a polypeptide that (i) is fewer than about 450 amino acid residues in length, (ii) includes an
5 amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, and (iii) exhibits β -secretase activity, as evidenced by an ability to cleave a substrate selected from the group consisting of the 695 amino acid isotype
15 of beta amyloid precursor protein (BAPP) between amino acids 596 and 597 thereof, MBP-C125wt and MBP-C125sw. Peptides which fit these criteria include, but are not limited to polypeptides which include the sequence SEQ ID NO: 75 [63-423], such as SEQ ID NO: 58
20 [46-452], SEQ ID NO: 58 [46-452], SEQ ID NO: 58 [46-452], SEQ ID NO: 74 [22-452], and may also include conservative substitutions within such sequences.

25 According to a further embodiment, the invention includes isolated protein compositions, such as those described above, in combination with a β -secretase substrate or
15 inhibitor molecule, such as MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof. Additional β -secretase-cleavable fragments useful in this regard are described in the specification hereof. Particularly useful inhibitors include peptides
30 derived from or including SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 72. Generally, such inhibitors will have K_i s of less than about 1 μ M. Such inhibitors may be labeled with a
20 detectable reporter molecule. Such labeled molecules are particularly useful, for example, in ligand binding assays.

35 In accordance with a further aspect, the invention includes protein compositions, such as those described above, expressed by a heterologous cell. In accordance with a further
40 embodiment, such cells may also co-express a β -secretase substrate or inhibitor protein or peptide. One or both of the expressed molecules may be heterologous to the cell.

45 In a related embodiment, the invention includes antibodies that bind specifically to a β -secretase protein comprising a polypeptide that includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof,
30 but which lacks significant immunoreactivity with a protein a sequence selected from the group consisting of SEQ ID NO: 2 [1-501] and SEQ ID NO: 43 [46-501].

50 In a further related embodiment, the invention includes isolated nucleic acids comprising a sequence of nucleotides that encodes a β -secretase protein that is at least 95%

5 identical to a protein selected from the group consisting of SEQ ID NO: 66 [22-501], SEQ ID
NO: 43[46-501], SEQ ID NO: 57 [1-419], SEQ ID NO: 74 [22-452], SEQ ID NO: 58 [46-
452], SEQ ID NO: 59 [1-452], SEQ ID NO: 60 [1-420], SEQ ID NO: 67 [58-501], SEQ ID
10 NO: 68 [58-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 70 [63-452], SEQ ID NO: 75 [63-
5 423], and SEQ ID NO: 71 [46-419], or a complementary sequence of any of such nucleotides.
Specifically excluded from this nucleotide is a nucleic acid encoding a protein having the
sequence SEQ ID NO: 2 [1-501].

15 Additionally, the invention includes an expression vector comprising such
isolated nucleic acids operably linked to the nucleic acid with regulatory sequences effective
20 for expression of the nucleic acid in a selected host cell, for heterologous expression. The
host cells can be a eukaryotic cell, a bacterial cell, an insect cell or a yeast cell. Such cells
can be used, for example, in a method of producing a recombinant β -secretase enzyme,
where the method further includes subjecting an extract or cultured medium from said cell to
25 an affinity matrix, such as a matrix formed from a β -secretase inhibitor molecule or antibody,
15 as detailed herein.

The invention is also directed to a method of screening for compounds that inhibit A β
production, comprising contacting a β -secretase polypeptide, such as those full-length or
30 truncated forms described above, with (i) a test compound and (ii) a β -secretase substrate,
and selecting the test compound as capable of inhibiting A β production if the β -secretase
20 polypeptide exhibits less β -secretase activity in the presence of than in the absence of the test
compound. Such an assay may be cell-based, with one or both of the enzyme and the
35 substrate produced by the cell, such as the co-expression cell referred to above. Kits
embodying such screening methods also form a part of the invention.

40 The screening method may further include administering a test compound to a
25 mammalian subject having Alzheimer's disease or Alzheimer's disease like pathology, and
selecting the compound as a therapeutic agent candidate if, following such administration, the
subject maintains or improves cognitive ability or the subject shows reduced plaque burden.
45 Preferably, such a subject is a comprising a transgene for human β -amyloid precursor protein
(β -APP), such as a mouse bearing a transgene which encodes a human β -APP, including a
30 mutant variants thereof, as exemplified in the specification.

5 In a related embodiment, the invention includes β -secretase inhibitor compound selected according to the methods described above. Such compounds may be selected, for example, from a phage display selection system ("library"), such as are known in
10 the art. According to another aspect, such libraries may be "biased" for the sequence peptide SEQ ID NO: 97 [P10-P4'D \rightarrow V]. Other inhibitors include, or may be derived from peptide inhibitors herein identified, such as inhibitors SEQ ID NO: 78, SEQ ID NO: 72, SEQ
15 ID NO: 78 and SEQ ID NO: 81.

Also forming part of the invention are knock-out mice, characterized by inactivation or deletion of an endogenous β -secretase gene, such as genes encodes a protein having at least
20 90% sequence identity to the sequence SEQ ID NO: 65. The deletion or inactivation may be inducible, such as by insertion of a Cre-lox expression system into the mouse genome.

According to a further related aspect, the invention includes a method of screening for drugs effective in the treatment of Alzheimer's disease or other cerebrovascular amyloidosis characterized by A β deposition. According to this aspect of the invention, a mammalian
25 subject characterized by overexpression of β -APP and/or deposition of A β is given a test compound selected for its ability to inhibit β -secretase activity a β -secretase protein according to claim 37. The compound is selected as a potential therapeutic drug compound, if it reduces the amount of A β deposition in said subject or if it maintains or improves cognitive ability in the subject. According to one preferred embodiment, the mammalian subject is a transgenic
30 mouse bearing a transgene encoding a human β -APP or a mutant thereof.

35 The invention also includes a method of treating a patient afflicted with or having a predilection for Alzheimer's disease or other cerebrovascular amyloidosis. According to this aspect, the enzymatic hydrolysis of APP to A β is blocked by administering to the patient a pharmaceutically effective dose of a compound effective to inhibit one or more of the various
40 forms of the enzyme described herein. According to another feature, the therapeutic compound is derived from a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 97. Such derivation may be effected by the various phage selection systems described herein, in conjunction with the screening
45 methods of the invention, or other such methods. Alternatively, or in addition, derivation may be achieved via rational chemistry approaches, including molecular modeling, known in
50 the medicinal chemistry art. Such compounds will preferably be rather potent inhibitors of β -secretase enzymatic activity, evidenced by a K_i of less than about 1-50 μ M in a MBP-

5 C125sw assay. Such compounds also form the basis for therapeutic drug compositions in accordance with the present invention, which may also include a pharmaceutically effective excipient.

10 According to yet another related aspect, the invention includes a
5 method of diagnosing the presence of or a predilection for Alzheimer's disease in a patient. This method includes detecting the expression level of a gene comprising a nucleic acid encoding β -secretase in a cell sample from said patient, and diagnosing the patient as having
15 or having a predilection for Alzheimer's disease, if said expression level is significantly greater than a pre-determined control expression level. Detectable nucleic acids, and primers
10 useful in such detection, are described in detail herein. Such nucleic acids may exclude a nucleic acid encoding the preproenzyme [1-501]. The invention is further directed to method
20 of diagnosing the presence of or a predilection for Alzheimer's disease in a patient, comprising measuring β -secretase enzymatic activity in a cell sample from said patient, and
25 diagnosing the patient as having or having a predilection for Alzheimer's disease, if said level
15 enzymatic activity level is significantly greater than a pre-determined control activity level. The diagnostic methods may be carried out in a whole cell assay and/or on a nucleic acid
derived from a cell sample of said patient.

30 The invention also includes a method of purifying a β -secretase protein enzyme molecule. According to this aspect, an impure sample containing β -secretase enzyme activity
20 with an affinity matrix which includes a β -secretase inhibitor, such as the various inhibitor molecules described herein.

35 These and other objects and features of the invention will become more fully apparent when the following detailed description of the invention is read in conjunction with the accompanying drawings.

40 25 **Brief Description of the Figures**

FIG. 1A shows the sequence of a polynucleotide (SEQ ID NO: 1) which encodes
45 human β -secretase translation product shown in FIG. 2A.

FIG. 1B shows the polynucleotide of FIG. 1A, including putative 5'- and 3'-
30 untranslated regions (SEQ ID NO: 44).

5 FIG. 2A shows the amino acid sequence (SEQ ID NO: 2)[1-501] of the predicted translation product of the open reading frame of the polynucleotide sequence shown in FIGS. 1A and 1B.

10 FIG. 2B shows the amino acid sequence of an active fragment of human β -secretase (SEQ ID NO: 43)[46-501].

15 FIG. 3A shows the translation product that encodes an active fragment of human β -secretase, 452stop, (amino acids 1-452 with reference to SEQ ID NO: 2; SEQ ID NO: 59) including a FLAG-epitope tag (underlined; SEQ ID NO: 45) at the C-terminus.

20 FIG. 3B shows the amino acid sequence of a fragment of human β -secretase (amino acids 46-452 (SEQ ID NO: 58) with reference to SEQ ID NO: 2; including a FLAG-epitope tag (underlined; SEQ ID NO: 45) at the C-terminus.

FIG. 4 shows an elution profile of recombinant β -secretase eluted from a gel filtration column.

25 FIG. 5 shows the full length amino acid sequence of β -secretase 1-501 (SEQ ID NO: 2), including the ORF which encodes it (SEQ ID NO: 1), with certain features indicated, such as "active-D" sites indicating the aspartic acid active catalytic sites, a transmembrane region commencing at position 453, as well as leader ("Signal") sequence (residues 1-22; SEQ ID NO: 46) and putative pro region (residues 23-45; SEQ ID NO: 47) and where the polynucleotide region corresponding the proenzyme region corresponding to amino acids 46-501 (SEQ ID NO: 43)(nt 135-1503) is shown as SEQ ID NO: 44.

30 FIGS. 6A and 6B show images of silver-stained SDS-PAGE gels on which purified β -secretase-containing fractions were run under reducing (6A) and non-reducing (6B) conditions.

35 FIG. 7 shows a silver-stained SDS-PAGE of β -secretase purified from heterologous 293T cells expressing the recombinant enzyme.

40 FIG. 8 shows a silver-stained SDS-PAGE of β -secretase purified from heterologous Cos A2 cells expressing the recombinant enzyme.

45 FIG. 9 shows a scheme in which primers derived from the polynucleotide (SEQ ID NO. 76 encoding N-terminus of purified naturally occurring β -secretase (SEQ ID NO. 77) were used to PCR-clone additional portions of the molecule, such as fragment SEQ ID NO. 79 encoding by nucleic acid SEQ ID NO. 78, as illustrated.

FIG. 10 shows an alignment of the amino acid sequence of human β -secretase ("Human Impain.seq," 1-501, SEQ ID NO: 2) compared to ("pBS/mImpain H#3 cons") consensus mouse sequence: SEQ ID NO: 65.

FIG. 11A shows the nucleotide sequence (SEQ ID NO: 80) of an insert used in preparing vector pCF.

FIG. 11B shows a linear schematic of pCEK.

FIG. 12 shows a schematic of pCEK.clone 27 used to transfect mammalian cells with β -secretase.

FIG. 13(A-E) shows the nucleotide sequence of pCEK clone 27 (SEQ ID NO: 48), with the OFR indicated by the amino acid sequence SEQ ID NO: 2.

FIG. 14A shows the a nucleotide sequence inserted into parent vector pCDNA3.

FIG. 14B shows a plot of β -secretase activity in cell lysates from COS cells transfected with vectors derived from clones encoding β -secretase.

FIGS. 15A shows an image of an SDS PAGE gel loaded with triplicate samples of the lysates made from heterologous cells transfected with mutant APP (751 wt) and β -galactosidase as control (lanes d) and from cells transfected with mutant APP (751 wt) and β -secretase (lanes f) where lanes a, b, and c show lysates from untreated cells, cells transfected with β -galactosidase alone and cells transfected with β -secretase alone, respectively, and lane e indicates markers.

FIG. 15B shows an image an image of an SDS PAGE gel loaded with triplicate samples of the lysates made from heterologous cells transfected with mutant APP (Swedish mutation) and β -galactosidase as control (lanes c) and from cells transfected with mutant APP (Swedish mutation) and β -secretase (lanes e) where lanes a and b show lysates from cells transfected with β -galactosidase alone and cells transfected with β -secretase alone, and lane d indicates markers.

FIGS. 16A and 16B show Western blots of cell supernatants tested for presence or increase in soluble APP (sAPP).

FIGS. 17A and 17B show Western blots of α -cleaved APP substrate in co-expression cells.

FIG. 18 shows A β (x-40) production in 293T cells cotransfected with APP and β -secretase.

FIG. 19A shows a schematic of an APP substrate fragment, and its use in conjunction with antibodies SW192 and 8E-192 in the assay.

FIG. 19B shows the β -secretase cleavage sites in the wild-type and Swedish APP sequence

FIG. 20 shows a schematic of a second APP substrate fragment derived from APP 638, and its use in conjunction with antibodies SW192 and 8E-192 in the assay.

FIG. 21 shows a schematic of pohCK751 vector.

Brief Description of the Sequences

This section briefly identifies the sequence identification numbers referred to herein. Number ranges shown in brackets here and throughout the specification are referenced to the amino acid sequence SEQ ID NO: 2, using conventional N \rightarrow C-terminus order.

SEQ ID NO: 1 is a nucleic acid sequence that encodes human β -secretase, including an active fragment, as exemplified herein.

SEQ ID NO: 2 is the predicted translation product of SEQ ID NO: 1 [1-501].

SEQ ID NOS: 3- 21 are degenerate oligonucleotide primers described in Example 1 (Table 4), designed from regions of SEQ ID NO: 2.

SEQ ID NOS: 22-41 are additional oligonucleotide primers used in PCR cloning methods described herein, shown in Table 5.

SEQ ID NO: 42 is a polynucleotide sequence that encodes the active enzyme β -secretase shown as SEQ ID NO: 43.

SEQ ID NO: 43 is the sequence of an active enzyme portion of human β -secretase, the N-terminus of which corresponds to the N-terminus of the predominant form of the protein isolated from natural sources [46-501].

SEQ ID NO: 44 is a polynucleotide which encodes SEQ ID NO: 2, including 5' and 3' untranslated regions.

SEQ ID NO: 45 is the FLAG sequence used in conjunction with certain polynucleotides.

SEQ ID NO: 46 is the putative leader region of β -secretase [1-22].

SEQ ID NO: 47 is the putative pre-pro region of β -secretase [23-45].

SEQ ID NO: 48 is the sequence of the clone pCEK Cl.27 (FIG. 13A-E).

5

SEQ ID NO: 49 is a nucleotide sequence of a fragment of the gene which encodes human β -secretase.

SEQ ID NO: 50 is the predicted translation product of SEQ ID NO: 49.

10

SEQ ID NO: 51 is the predicted internal amino acid sequence of a portion of human

5 β -secretase.

SEQ ID NOS: 52 and 53 are peptide substrates suitable for use in β -secretase assays used in the present invention.

15

SEQ ID NO: 54 is a peptide sequence cleavage site recognized by human β -secretase.

SEQ ID NO: 55 is amino acids 46-69 of SEQ ID NO: 2.

10

SEQ ID NO: 56 is an internal peptide just N-terminal to the transmembrane domain of β -secretase.

20

SEQ ID NO: 57 is β -secretase [1-419].

SEQ ID NO: 58 is β -secretase [46-452].

25

SEQ ID NO: 59 is β -secretase [1-452].

15

SEQ ID NO: 60 is β -secretase [1-420].

SEQ ID NO: 61 is EVM[hydroxyethylene]AEF.

SEQ ID NO: 62 is the amino acid sequence of the transmembrane domain of β -secretase shown in (FIG. 5).

30

SEQ ID NO: 63 is P26-P4' of APPwt.

20

SEQ ID NO: 64 is P26-P1' of APPwt.

35

SEQ ID NO: 65 is mouse β -secretase (FIG. 10, lower sequence).

SEQ ID NO: 66 is β -secretase [22-501].

SEQ ID NO: 67 is β -secretase [58-501].

40

SEQ ID NO: 68 is β -secretase [58-452].

25

SEQ ID NO: 69 is β -secretase [63-501].

SEQ ID NO: 70 is β -secretase [63-452].

SEQ ID NO: 71 is β -secretase [46-419].

45

SEQ ID NO: 72 is P10-P4'staD \rightarrow V.

SEQ ID NO: 73 is P4-P4'staD \rightarrow V (KTEEISEVN[sta]VAEF).

30

SEQ ID NO: 74 is β -secretase [22-452].

50

SEQ ID NO: 75 is β -secretase [63-423].

55

5

SEQ ID NO: 76 is nucleic acid encoding the N-terminus of naturally occurring β -secretase.

10

SEQ ID NO: 77 is a peptide fragment at the N-terminus of naturally occurring β -secretase.

5

SEQ ID NO: 78 is a P3-P4'XD \rightarrow V (VMXVAEF, where X is hydroxyethylene or statine).

15

SEQ ID NO: 79 is a peptide fragment of naturally occurring occurring β -secretase.

SEQ ID NO: 80 is a nucleotide insert in vector pCF used herein.

10

SEQ ID NO: 81 is P4-P4'XD \rightarrow V (EVMXVAEF, where X is hydroxyethylene or statine).

20

SEQ ID NO: 82 is APP fragment SEVKMDAEF (P5-P4'wt).

SEQ ID NO: 83 is APP fragment SEVNLDAEF (P5-P4'sw).

SEQ ID NO: 84 is APP fragment SEVKLDAEF.

25

SEQ ID NO: 85 is APP fragment SEVKFDAEF.

15

SEQ ID NO: 86 is APP fragment SEVNFDAEF.

SEQ ID NO: 87 is APP fragment SEVKMAAEF.

SEQ ID NO: 88 is APP fragment SEVNLADEF.

30

SEQ ID NO: 89 is APP fragment SEVKLAAEF.

SEQ ID NO: 90 is APP fragment SEVKMLAEF.

20

SEQ ID NO: 91 is APP fragment SEVNLLAEF.

SEQ ID NO: 92 is APP fragment SEVKLLAEF.

35

SEQ ID NO: 93 is APP fragment SEVKFAAEF.

SEQ ID NO: 94 is APP fragment SEVNFAAEF.

SEQ ID NO: 95 is APP fragment SEVKFLAEF.

40

25

SEQ ID NO: 96 is APP fragment SEVNFLAEF.

SEQ ID NO: 97 is APP-derived fragment P10-P4'(D \rightarrow V): KTEEISEVNLVAEF

Detailed Description of the Invention

45

50

55

5
I. Definitions

Unless otherwise indicated, all terms used herein have the same meaning as they would to one skilled in the art of the present invention. Practitioners are particularly directed to
10 Sambrook, *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Second Edition), Cold
5 Spring Harbor Press, Plainview, N.Y., and Ausubel, F.M., *et al.* (1998) *Current Protocols in*
Molecular Biology, John Wiley & Sons, New York, NY, for definitions, terms of art and
15 standard methods known in the art of molecular biology, particularly as it relates to the cloning
protocols described herein. It is understood that this invention is not limited to the particular
methodology, protocols, and reagents described, as these may be varied to produce the same
10 result.

20 The terms "polynucleotide" and "nucleic acid" are used interchangeably herein and refer
to a polymeric molecule having a backbone that supports bases capable of hydrogen bonding to
typical polynucleotides, where the polymer backbone presents the bases in a manner to permit
such hydrogen bonding in a sequence specific fashion between the polymeric molecule and a
25 typical polynucleotide (*e.g.*, single-stranded DNA). Such bases are typically inosine, adenosine,
15 guanosine, cytosine, uracil and thymidine. Polymeric molecules include double and single
stranded RNA and DNA, and backbone modifications thereof, for example, methylphosphonate
linkages.
30

The term "vector" refers to a polynucleotide having a nucleotide sequence that can
20 assimilate new nucleic acids, and propagate those new sequences in an appropriate host.
Vectors include, but are not limited to recombinant plasmids and viruses. The vector (*e.g.*,
35 plasmid or recombinant virus) comprising the nucleic acid of the invention can be in a carrier,
for example, a plasmid complexed to protein, a plasmid complexed with lipid-based nucleic
acid transduction systems, or other non-viral carrier systems.

40 25 The term "polypeptide" as used herein refers to a compound made up of a single chain
of amino acid residues linked by peptide bonds. The term "protein" may be synonymous with
the term "polypeptide" or may refer to a complex of two or more polypeptides.

The term "modified", when referring to a polypeptide of the invention, means a
45 polypeptide which is modified either by natural processes, such as processing or other post-
translational modifications, or by chemical modification techniques which are well known in
30 the art. Among the numerous known modifications which may be present include, but are not
50 limited to, acetylation, acylation, amidation, ADP-ribosylation, glycosylation, GPI anchor

5 formation, covalent attachment of a lipid or lipid derivative, methylation, myristylation, pegylation, prenylation, phosphorylation, ubiquitination, or any similar process.

The term "β-secretase" is defined in Section III, herein.

10 The term "biologically active" used in conjunction with the term β-secretase refers to possession of a β-secretase enzyme activity, such as the ability to cleave β-amyloid precursor protein (APP) to produce β-amyloid peptide (Aβ).

15 The term "fragment," when referring to β-secretase of the invention, means a polypeptide which has an amino acid sequence which is the same as part of but not all of the amino acid sequence of full-length β-secretase polypeptide. In the context of the present invention, the full length β-secretase is generally identified as SEQ ID NO: 2, the ORF of the full-length nucleotide; however, according to a discovery of the invention, the naturally occurring active form is probably one or more N-terminal truncated versions, such as amino acids 46-501, 22-501, 58-501 or 63-501; other active forms are C-terminal truncated forms ending between about amino acids 450 and 452. The numbering system used throughout is based on the numbering of the sequence SEQ ID NO: 2.

20 An "active fragment" is a β-secretase fragment that retains at least one of the functions or activities of β-secretase, including but not limited to the β-secretase enzyme activity discussed above and/or ability to bind to the inhibitor substrate described herein as P10-P4'staD->V. Fragments contemplated include, but are not limited to, a β-secretase fragment which retains the ability to cleave β-amyloid precursor protein to produce β-amyloid peptide. Such a fragment preferably includes at least 350, and more preferably at least 400, contiguous amino acids or conservative substitutions thereof of β-secretase, as described herein. More preferably, the fragment includes active aspartyl acid residues in the structural proximities identified and defined by the primary polypeptide structure shown as SEQ ID NO: 2 and also denoted as "Active-D" sites herein.

25 A "conservative substitution" refers to the substitution of an amino acid in one class by an amino acid in the same class, where a class is defined by common physicochemical amino acid sidechain properties and high substitution frequencies in homologous proteins found in nature (as determined, e.g., by a standard Dayhoff frequency exchange matrix or BLOSUM matrix). Six general classes of amino acid sidechains, categorized as described above, include: Class I (Cys); Class II (Ser, Thr, Pro, Ala, Gly); Class III (Asn, Asp, Gln, Glu); Class IV (His, Arg, Lys); Class V (Ile, Leu, Val, Met); and Class VI (Phe, Tyr, Trp).

5 For example, substitution of an Asp for another class III residue such as Asn, Gln, or Glu, is considered to be a conservative substitution.

10 "Optimal alignment" is defined as an alignment giving the highest percent identity score. Such alignment can be performed using a variety of commercially available sequence analysis programs, such as the local alignment program LALIGN using a ktp of 1, default parameters and the default PAM. A preferred alignment is the pairwise alignment using the CLUSTAL-W program in MacVector, operated with default parameters, including an open gap penalty of 10.0, an extended gap penalty of 0.1, and a BLOSUM30 similarity matrix.

15 "Percent sequence identity," with respect to two amino acid or polynucleotide sequences, refers to the percentage of residues that are identical in the two sequences when the sequences are optimally aligned. Thus, 80% amino acid sequence identity means that 80% of the amino acids in two or more optimally aligned polypeptide sequences are identical. If a gap needs to be inserted into a first sequence to optimally align it with a second sequence, the percent identity is calculated using only the residues that are paired with a corresponding amino acid residue (i.e., the calculation does not consider residues in the second sequences that are in the "gap" of the first sequence.

20 A first polypeptide region is said to "correspond" to a second polypeptide region when the regions are essentially co-extensive when the sequences containing the regions are aligned using a sequence alignment program, as above. Corresponding polypeptide regions typically contain a similar, if not identical, number of residues. It will be understood, however, that corresponding regions may contain insertions or deletions of residues with respect to one another, as well as some differences in their sequences.

25 A first polynucleotide region is said to "correspond" to a second polynucleotide region when the regions are essentially co-extensive when the sequences containing the regions are aligned using a sequence alignment program, as above. Corresponding polynucleotide regions typically contain a similar, if not identical, number of residues. It will be understood, however, that corresponding regions may contain insertions or deletions of bases with respect to one another, as well as some differences in their sequences.

30 The term "sequence identity" means nucleic acid or amino acid sequence identity in two or more aligned sequences, aligned as defined above.

35 "Sequence similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. Thus, 80% protein sequence similarity means that 80% of

5 the amino acid residues in two or more aligned protein sequences are conserved amino acid residues, *i.e.* are conservative substitutions.

10 "Hybridization" includes any process by which a strand of a nucleic acid joins with a complementary nucleic acid strand through base pairing. Thus, strictly speaking, the term
5 refers to the ability of the complement of the target sequence to bind to the test sequence, or vice-versa.

15 "Hybridization conditions" are based in part on the melting temperature (T_m) of the nucleic acid binding complex or probe and are typically classified by degree of "stringency" of the conditions under which hybridization is measured. The specific conditions that define
10 various degrees of stringency (*i.e.*, high, medium, low) depend on the nature of the polynucleotide to which hybridization is desired, particularly its percent GC content, and can
20 be determined empirically according to methods known in the art. Functionally, maximum stringency conditions may be used to identify nucleic acid sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to
25 identify nucleic acid sequences having about 80% or more sequence identity with the probe.

30 The term "gene" as used herein means the segment of DNA involved in producing a polypeptide chain; it may include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

35 The term "isolated" means that the material is removed from its original environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same
40 polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such isolated polynucleotides may be part of a vector and/or such
25 polynucleotides or polypeptides may be part of a composition, such as a recombinantly produced cell (heterologous cell) expressing the polypeptide, and still be isolated in that such vector or composition is not part of its natural environment.

45 An "isolated polynucleotide having a sequence which encodes β -secretase" is a polynucleotide that contains the coding sequence of β -secretase, or an active fragment
30 thereof, (i) alone, (ii) in combination with additional coding sequences, such as fusion protein or signal peptide, in which the β -secretase coding sequence is the dominant coding
50 sequence, (iii) in combination with non-coding sequences, such as introns and control elements, such as promoter and terminator elements or 5' and/or 3' untranslated regions,

5 effective for expression of the coding sequence in a suitable host, and/or (iv) in a vector or host environment in which the β -secretase coding sequence is a heterologous gene.

10 The terms "heterologous DNA," "heterologous RNA," "heterologous nucleic acid," "heterologous gene," and "heterologous polynucleotide" refer to nucleotides that are not
5 endogenous to the cell or part of the genome in which they are present; generally such nucleotides have been added to the cell, by transfection, microinjection, electroporation, or the like. Such nucleotides generally include at least one coding sequence, but this coding
15 sequence need not be expressed.

The term "heterologous cell" refers to a recombinantly produced cell that contains at
10 least one heterologous DNA molecule.

20 A "recombinant protein" is a protein isolated, purified, or identified by virtue of expression in a heterologous cell, said cell having been transduced or transfected, either transiently or stably, with a recombinant expression vector engineered to drive expression of the protein in the host cell.

25 15 The term "expression" means that a protein is produced by a cell, usually as a result of transfection of the cell with a heterologous nucleic acid.

30 "Co-expression" is a process by which two or more proteins or RNA species of interest are expressed in a single cell. Co-expression of the two or more proteins is typically achieved by transfection of the cell with one or more recombinant expression vectors(s) that
20 carry coding sequences for the proteins. In the context of the present invention, for example, a cell can be said to "co-express" two proteins, if one or both of the proteins is heterologous
35 to the cell.

40 25 The term "expression vector" refers to vectors that have the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

45 The terms "purified" or "substantially purified" refer to molecules, either polynucleotides or polypeptides, that are removed from their natural environment, isolated or separated, and are at least 90% and more preferably at least 95-99% free from other
30 components with which they are naturally associated. The foregoing notwithstanding, such a descriptor does not preclude the presence in the same sample of splice- or other protein
50 variants (glycosylation variants) in the same, otherwise homogeneous, sample.

5 A protein or polypeptide is generally considered to be "purified to apparent homogeneity" if a sample containing it shows a single protein band on a silver-stained polyacrylamide electrophoretic gel.

10 The term "crystallized protein" means a protein that has co-precipitated out of solution in pure crystals consisting only of the crystal, but possibly including other components that are tightly bound to the protein.

15 A "variant" polynucleotide sequence may encode a "variant" amino acid sequence that is altered by one or more amino acids from the reference polypeptide sequence. The variant polynucleotide sequence may encode a variant amino acid sequence, which contains
20 "conservative" substitutions, wherein the substituted amino acid has structural or chemical properties similar to the amino acid which it replaces. In addition, or alternatively, the variant polynucleotide sequence may encode a variant amino acid sequence, which contains "non-conservative" substitutions, wherein the substituted amino acid has dissimilar structural or chemical properties to the amino acid which it replaces. Variant polynucleotides may also
25 encode variant amino acid sequences, which contain amino acid insertions or deletions, or both. Furthermore, a variant polynucleotide may encode the same polypeptide as the reference polynucleotide sequence but, due to the degeneracy of the genetic code, has a polynucleotide sequence that is altered by one or more bases from the reference polynucleotide sequence.

30 An "allelic variant" is an alternate form of a polynucleotide sequence, which may have a substitution, deletion or addition of one or more nucleotides that does not substantially alter the function of the encoded polypeptide.

35 "Alternative splicing" is a process whereby multiple polypeptide isoforms are generated from a single gene, and involves the splicing together of nonconsecutive exons during the processing of some, but not all, transcripts of the gene. Thus, a particular exon
40 may be connected to any one of several alternative exons to form messenger RNAs. The alternatively-spliced mRNAs produce polypeptides ("splice variants") in which some parts are common while other parts are different.

45 "Splice variants" of β -secretase, when referred to in the context of an mRNA transcript, are mRNAs produced by alternative splicing of coding regions, i.e., exons, from the β -secretase gene.

5 "Splice variants" of β -secretase, when referred to in the context of the protein itself, are β -secretase translation products that are encoded by alternatively-spliced β -secretase mRNA transcripts.

10 A "mutant" amino acid or polynucleotide sequence is a variant amino acid sequence, or a variant polynucleotide sequence, which encodes a variant amino acid sequence that has significantly altered biological activity or function from that of the naturally occurring protein.

15 A "substitution" results from the replacement of one or more nucleotides or amino acids by different nucleotides or amino acids, respectively.

20 The term "modulate" as used herein refers to the change in activity of the polypeptide of the invention. Modulation may relate to an increase or a decrease in biological activity, binding characteristics, or any other biological, functional, or immunological property of the molecule.

25 The terms "antagonist" and "inhibitor" are used interchangeably herein and refer to a molecule which, when bound to the polypeptide of the present invention, modulates the activity of enzyme by blocking, decreasing, or shortening the duration of the biological activity. An antagonist as used herein may also be referred to as a " β -secretase inhibitor" or " β -secretase blocker." Antagonists may themselves be polypeptides, nucleic acids, carbohydrates, lipids, small molecules (usually less than 1000 kD), or derivatives thereof, or any other ligand which binds to and modulates the activity of the enzyme.

35 β -Secretase Compositions

40 The present invention provides an isolated, active human β -secretase enzyme, which is further characterized as an aspartyl (aspartic) protease or proteinase, optionally, in purified form. As defined more fully in the sections that follow, β -secretase exhibits a proteolytic activity that is involved in the generation of β -amyloid peptide from β -amyloid precursor protein (APP), such as is described in U.S. Patent 5,744,346, incorporated herein by reference. Alternatively, or in addition, the β -secretase is characterized by its ability to bind, with moderately high affinity, to an inhibitor substrate described herein as P10-P4' staD \rightarrow V (SEQ ID NO.: 72). According to an important feature of the present invention, a human form of β -secretase has been isolated, and its naturally occurring form has been characterized, purified and sequenced.

5 According to another aspect of the invention, nucleotide sequences encoding the enzyme have been identified. In addition, the enzyme has been further modified for expression in altered forms, such as truncated forms, which have similar protease activity to
10 the naturally occurring or full length recombinant enzyme. Using the information provided herein, practitioners can isolate DNA encoding various active forms of the protein from
15 available sources and can express the protein recombinantly in a convenient expression system. Alternatively and in addition, practitioners can purify the enzyme from natural or recombinant sources and use it in purified form to further characterize its structure and function. According to a further feature of the invention, polynucleotides and proteins of the
20 invention are particularly useful in a variety of screening assay formats, including cell-based screening for drugs that inhibit the enzyme. Examples of uses of such assays, as well as additional utilities for the compositions are provided in Section IV, below.

25 β -secretase is of particular interest due to its activity and involvement in generating fibril peptide components that are the major components of amyloid plaques in the central nervous system (CNS), such as are seen in Alzheimer's disease, Down's syndrome and other
15 CNS disorders. Accordingly, a useful feature of the present invention includes an isolated form of the enzyme that can be used, for example, to screen for inhibitory substances which are candidates for therapeutics for such disorders.

30 A. Isolation of Polynucleotides encoding Human β -secretase Polynucleotides encoding human β -secretase were obtained by PCR cloning and hybridization techniques as
20 detailed in Examples 1-3 and described below. FIG. 1A shows the sequence of a polynucleotide (SEQ ID NO: 1) which encodes a form of human β -secretase (SEQ ID NO: 2 [1-501]). Polynucleotides encoding human β -secretase are conveniently isolated from any of a
35 number of human tissues, preferably tissues of neuronal origin, including but not limited to neuronal cell lines such as the commercially available human neuroblastoma cell line IMR-32
40 available from the American Type Culture Collection (Manassas, VA; ATTC CCL 127) and human fetal brain, such as a human fetal brain cDNA library available from OriGene Technologies, Inc. (Rockville, MD).

45 Briefly, human β -secretase coding regions were isolated by methods well known in the art, using hybridization probes derived from the coding sequence provided as SEQ ID
30 NO: 1. Such probes can be designed and made by methods well known in the art. Exemplary probes, including degenerate probes, are described in Example 1. Alternatively, a
50

5 cDNA library is screened by PCR, using, for example, the primers and conditions described in Example 2 herein. Such methods are discussed in more detail in Part B, below.

10 cDNA libraries were also screened using a 3'-RACE (Rapid Amplification of cDNA Ends) protocol according to methods well known in the art (White, B.A., ed., PCR Cloning
5 Protocols; Humana Press, Totowa, NJ, 1997; shown schematically in FIG. 9). Here primers derived from the 5' portion of SEQ ID NO: 1 are added to partial cDNA substrate clone found by screening a fetal brain cDNA library as described above. A representative 3'RACE
15 reaction used in determining the longer sequence is detailed in Example 3 and is described in more detail in Part B, below.

20 Human β -secretase, as well as additional members of the neuronal aspartyl protease family described herein may be identified by the use of random degenerate primers designed in accordance with any portion of the polypeptide sequence shown as SEQ ID NO: 2. For
25 example, in experiments carried out in support of the present invention, and detailed in Example 1 herein, eight degenerate primer pools, each 8-fold degenerate, were designed
15 based on a unique 22 amino acid peptide region selected from SEQ ID: 2. Such techniques can be used to identify further similar sequences from other species and/or representing other members of this protease family.

30 Preparation of polynucleotides

20 The polynucleotides described herein may be obtained by screening cDNA libraries using oligonucleotide probes, which can hybridize to and/or PCR-amplify polynucleotides
35 that encode human β -secretase, as disclosed above. cDNA libraries prepared from a variety of tissues are commercially available, and procedures for screening and isolating cDNA clones are well known to those of skill in the art. Genomic libraries can likewise be screened
40 25 to obtain genomic sequences including regulatory regions and introns. Such techniques are described in, for example, Sambrook *et al.* (1989) Molecular Cloning: A Laboratory Manual (2nd Edition), Cold Spring Harbor Press, Plainview, N.Y. and Ausubel, FM *et al.* (1998) Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y.

45 The polynucleotides may be extended to obtain upstream and downstream sequences
30 such as promoters, regulatory elements, and 5' and 3' untranslated regions (UTRs). Extension of the available transcript sequence may be performed by numerous methods known to those of skill in the art, such as PCR or primer extension (Sambrook *et al.*, supra), or by the RACE
50

5 method using, for example, the MARATHON RACE kit (Cat. # K1802-1; Clontech, Palo Alto, CA).

10 Alternatively, the technique of "restriction-site" PCR (Gobinda *et al.* (1993) PCR Methods Applic. 2:318-22), which uses universal primers to retrieve flanking sequence
5 adjacent a known locus, may be employed to generate additional coding regions. First, genomic DNA is amplified in the presence of primer to a linker sequence and a primer specific to the known region. The amplified sequences are subjected to a second round of
15 PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and
10 sequenced using reverse transcriptase.

20 Inverse PCR can be used to amplify or extend sequences using divergent primers based on a known region (Triglia T *et al.* (1988) Nucleic Acids Res 16:8186). The primers may be designed using OLIGO(R) 4.06 Primer Analysis Software (1992; National Biosciences Inc, Plymouth, Minn.), or another appropriate program, to be 22-30 nucleotides
25 in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by
30 intramolecular ligation and used as a PCR template.

35 Capture PCR (Lagerstrom M *et al.* (1991) PCR Methods Applic 1:111-19) is a
20 method for PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. Capture PCR also requires multiple restriction enzyme digestions and ligations to place an engineered double-stranded sequence into a
35 flanking part of the DNA molecule before PCR.

40 Another method which may be used to retrieve flanking sequences is that of Parker, JD *et al.* (1991; Nucleic Acids Res 19:3055-60). Additionally, one can use PCR, nested
25 primers and PromoterFinder(TM) libraries to "walk in" genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions. Preferred libraries for screening for full length cDNAs are ones that have been
45 size-selected to include larger cDNAs. Also, random primed libraries are preferred in that
30 they will contain more sequences which contain the 5' and upstream regions of genes. A randomly primed library may be particularly useful if an oligo d(T) library does not yield a
50 full-length cDNA. Genomic libraries are useful for extension into the 5' nontranslated regulatory region.

5 The polynucleotides and oligonucleotides of the invention can also be prepared by solid-phase methods, according to known synthetic methods. Typically, fragments of up to about 100 bases are individually synthesized, then joined to form continuous sequences up to
10 several hundred bases.

5 B. Isolation of β -Secretase

The amino acid sequence for a full-length human β -secretase translation product is shown as SEQ ID NO: 2 in FIG. 2A. According to the discovery of the present invention,
15 this sequence represents a "pre pro" form of the enzyme that was deduced from the nucleotide sequence information described in the previous section in conjunction with the methods described below. Comparison of this sequence with sequences determined from the
20 biologically active form of the enzyme purified from natural sources, as described in Part 4, below, indicate that it is likely that an active and predominant form of the enzyme is represented by sequence shown in FIG. 2B (SEQ ID NO: 43), in which the first 45 amino acids of the open-reading frame deduced sequence have been removed. This suggests that
25 the enzyme may be post-translationally modified by proteolytic activity, which may be autocatalytic in nature. Further analysis, illustrated by the schematics shown in FIG. 5 herein, indicates that the enzyme contains a hydrophobic, putative transmembrane region near its C-terminus. As described below, a further discovery of the present invention is that the
30 enzyme can be truncated prior to this transmembrane region and still retain β -secretase activity.

35 1. Purification of β -secretase from Natural and Recombinant Sources

According to an important feature of the present invention, β -secretase has now been purified from natural and recombinant sources. U.S. Patent 5,744,346, incorporated herein by reference, describes isolation of β -secretase in a single peak having an apparent molecular
40 25 weight of 260-300,000 (Daltons) by gel exclusion chromatography. It is a discovery of the present invention that the native enzyme can be purified to apparent homogeneity by affinity column chromatography. The methods revealed herein have been used on preparations from brain tissue as well as on preparations from 293T and recombinant cells; accordingly, these
45 methods are believed to be generally applicable over a variety of tissue sources. The practitioner will realize that certain of the preparation steps, particularly the initial steps, may require modification to accommodate a particular tissue source and will adapt such
30 procedures according to methods known in the art. Methods for purifying β -secretase from
50

human brain as well as from cells are detailed in Example 5. Briefly, cell membranes or brain tissue are homogenized, fractionated, and subjected to various types of column chromatographic matrices, including wheat germ agglutinin-agarose (WGA), anion exchange chromatography and size exclusion. Activity of fractions can be measured using any appropriate assay for β -secretase activity, such as the MBP-C125 cleavage assay detailed in Example 4. Fractions containing β -secretase activity elute from this column in a peak elution volume corresponding to a size of about 260-300 kilodaltons.

The foregoing purification scheme, which yields approximately 1,500-fold purification, is similar to that described in detail in U.S. Patent 5,744,346, incorporated herein by reference. In accordance with the present invention, further purification can be achieved by applying the cation exchange flow-through material to an affinity column that employs as its affinity matrix a specific inhibitor of β -secretase, termed "P10—P4'staD->V" (NH₂-KTEEISEVN[sta]VAEF-CO₂H; SEQ ID NO.: 72). This inhibitor, and methods for making a Sepharose affinity column which incorporates it, are described in Example 7. After washing the column, β -secretase and a limited number of contaminating proteins were eluted with pH 9.5 borate buffer. The eluate was then fractionated by anion exchange HPLC, using a Mini-Q column. Fractions containing the activity peak were pooled to give the final β -secretase preparation. Results of an exemplary run using this purification scheme are summarized in Table 1. FIG. 6A shows a picture of a silver-stained SDS PAGE gel run under reducing conditions, in which β -secretase runs as a 70 kilodalton band. The same fractions run under non-reducing conditions (FIG. 6B) provide evidence for disulfide cross-linked oligomers. When the anion exchange pool fractions 18-21 (see FIG. 6B) were treated with dithiothreitol (DTT) and re-chromatographed on a Mini Q column, then subjected to SDS-PAGE under non-reducing conditions, a single band running at about 70 kilodaltons was observed. Surprisingly, the purity of this preparation is at least about 200 fold higher than the previously purified material, described in U.S. Patent 5,744,346. By way of comparison, the most pure fraction described therein exhibited a specific activity of about 253 nM/h/ μ g protein, taking into consideration the MW of substrate MBP-C26sw (45 kilodaltons). The present method therefore provides a preparation that is at least about 1000-fold higher purity (affinity eluate) and as high as about 6000-fold higher purity than that preparation, which represented at least 5 to 100-fold higher purity than the enzyme present in a solubilized but unenriched membrane fraction from human 293 cells.

Table 1

Preparation of β -secretase from Human Brain

	Total Activity ^a nM/h	Specific Activity ^b nM/h/ μ g prot.	% Yield	Purification (fold)
Brain Extract	19,311,150	4.7	100	1
WGA Eluate	21,189,600	81.4	110	17
Affinity Eluate	11,175,000	257,500	53	54,837
Anion Exchange Pool	3,267,685	1,485,312	17	316,309

^aActivity in MBP-C125sw assay^bSpecific Activity = $\frac{(\text{Product conc. nM})(\text{Dilution factor})}{(\text{Enzyme sol. vol})(\text{Incub. time h})(\text{Enzyme conc. } \mu\text{g/vol})}$

Example 5 also describes purification schemes used for purifying recombinant materials from heterologous cells transfected with the β -secretase coding sequence. Results from these purifications are illustrated in FIGS. 7 and 8. Further experiments carried out in support of the present invention, showed that the recombinant material has an apparent molecular weight in the range from 260,000 to 300,000 Daltons when measured by gel exclusion chromatography. FIG. 4 shows an activity profile of this preparation run on a gel exclusion chromatography column, such as a Superdex 200 (26/60) column, according to the methods described in U.S. Patent 5,744,346, incorporated herein by reference.

1. Sequencing of β -secretase Protein

A schematic overview summarizing methods and results for determining the cDNA sequence encoding the N-terminal peptide sequence determined from purified β -secretase is shown in FIG. 9. N-terminal sequencing of purified β -secretase protein isolated from natural sources yielded a 21-residue peptide sequence, SEQ ID NO. 77, as described above. This peptide sequence, and its reverse translated fully degenerate nucleotide sequence, SEQ ID NO. 76, is shown in the top portion of FIG. 4. Two partially degenerate primer sets used for RT-PCR amplification of a cDNA fragment encoding this peptide are also summarized in FIG. 4. Primer set 1 consisted of DNA nucleotide primers #3427-3434, shown in Table 3 (Example 3). Matrix RT-PCR using combinations of primers from this set with cDNA reverse transcribed from primary human neuronal cultures as template yielded the predicted 54 bp cDNA product with primers #3428 – 3433, also described in Table 3.

In further experiments carried out in support of the present invention, it was found that oligonucleotides from primer sets 1 and 2 could also be used to amplify cDNA fragments of the predicted size from mouse brain mRNA. DNA sequence demonstrated that such primers could also be used to clone the murine homolog(s) and other species homologs of

human β -secretase and/or additional members of the aspartyl protease family described herein by standard RACE-PCR technology. The sequence of a murine homolog is presented in FIG. 10 (lower sequence; "pBS/MuImPain H#3 cons"); SEQ ID NO. 65. The murine polypeptide sequence is about 95% identical to the human polypeptide sequence.

2. 5' and 3' RACE-PCR for Additional Sequence, Cloning, and mRNA Analysis

The unambiguous internal nucleotide sequence from the amplified fragment provided information which facilitated the design of internal primers matching the upper (coding) strand for 3' RACE, and lower (non-coding) strand for 5' RACE (Frohman, M. A., M. K. Dush and G. R. Martin (1988). "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene specific oligo-nucleotide primer." Proc. Natl. Acad. Sci. U.S.A. 85(23): 8998-9002.) The DNA primers used for this experiment (#3459 & #3460) are illustrated schematically in FIG. 9, and the exact sequence of these primers is presented in Table 4 of Example 3.

Primers #3459 and #3476 (Table 5) were used for initial 3' RACE amplification of downstream sequences from the IMR-32 cDNA library in the vector pLPCXlox. The library had previously been sub-divided into 100 pools of 5,000 clones per pool, and plasmid DNA was isolated from each pool. A survey of the 100 pools with the primers described in Part 2, above, identified individual pools containing β -secretase clones from the library. Such clones can be used for RACE-PCR analysis.

An approximately 1.8 Kb PCR fragment was observed by agarose gel fractionation of the reaction products. The PCR product was purified from the gel and subjected to DNA sequence analysis using primer #3459 (Table 5). The resulting clone sequence, designated 23A, was determined. Six of the first seven deduced amino-acids from one of the reading frames of 23A were an exact match with the last 7 amino-acids of the N-terminal sequence (SEQ ID NO. 77) determined from the purified protein isolated from natural sources in other experiments carried out in support of this invention. This observation provided internal validation of the sequences, and defined the proper reading frame downstream. Furthermore, this DNA sequence facilitated design of additional primers for extending the sequence further downstream, verifying the sequence by sequencing the opposite strand in the upstream direction, and further facilitated isolating the cDNA clone.

A DNA sequence of human β -secretase is illustrated as SEQ ID NO: 42 corresponding to SEQ ID NO: 1 including 5'- and 3'-untranslated regions. This sequence was determined from a partial cDNA clone (9C7e.35) isolated from a commercially available

5 human fetal brain cDNA library purchased from OriGene™, the 3' RACE product 23A, and additional clones – a total of 12 independent cDNA clones were used to determine the composite sequence. The composite sequence was assembled by sequencing overlapping
10 stretches of DNA from both strands of the clone or PCR fragment. The predicted full length translation product is shown as SEQ ID NO: 2 in Fig. 1B.

4. Tissue Distribution of β -secretase and Related Transcripts

15 Oligonucleotide primer #3460 (SEQ ID NO. 39, Table 5) was employed as an end-labeled probe on Northern blots to determine the size of the transcript encoding β -secretase and to examine its expression in IMR-32 cells. Additional primers were used to isolate the
20 mouse cDNA and to characterize mouse tissues, using Marathon RACE ready cDNA preparations (Clontech, Palo Alto, CA). TABLE 2 summarizes the results of experiments in which various human and murine tissues were tested for the presence of β -secretase-encoding transcripts by PCR or Northern blotting.

25 For example, the oligo-nucleotide probe 3460 (SEQ ID NO: 39) hybridized to a 2 Kb transcript in IMR-32 cells, indicating that the mRNA encoding the β -secretase enzyme is 2 Kb in size in this tissue. Northern blot analysis of total RNA isolated from the human T-cell
30 line Jurkat, and human myelomonocyte line Thp1 with the 3460 oligo-nucleotide probe 3460 also revealed the presence of a 2 kb transcript in these cells.

35 The oligonucleotide probe #3460 also hybridizes to a ~2 kb transcript in Northern blots containing RNA from all human organs examined to date, from both adult and fetal tissue. The organs surveyed include heart, brain, liver, pancreas, placenta, lung, muscle,
40 uterus, bladder, kidney, spleen, skin, and small intestine. In addition, certain tissues, e.g. pancreas, liver, brain, muscle, uterus, bladder, kidney, spleen and lung, show expression of larger transcripts of ~4.5 kb, 5 kb, and 6.5 kb which hybridize with oligonucleotide probe
45 #3460.

50 In further experiments carried out in support of the present invention, Northern blot results were obtained with oligonucleotide probe #3460 by employing a riboprobe derived from SEQ ID NO: 1, encompassing nucleotides #155-1014. This clone provides an 860 bp
55 riboprobe, encompassing the catalytic domain-encoding portion of β -secretase, for high stringency hybridization. This probe hybridized with high specificity to the exact match mRNA expressed in the samples being examined. Northern blots of mRNA isolated from IMR-32 and 1^{HNC} probed with this riboprobe revealed the presence of the 2 kb transcript

5 previously detected with oligonucleotide #3460, as well as a novel, higher MW transcript of
~5 kb. Hybridization of RNA from adult and fetal human tissues with this 860 nt riboprobe
also confirmed the result obtained with the oligonucleotide probe #3460. The mRNA
10 encoding β -secretase is expressed in all tissues examined, predominantly as an ~5 kb
5 transcript. In adult, its expression appeared lowest in brain, placenta, and lung, intermediate
in uterus, and bladder, and highest in heart, liver, pancreas, muscle, kidney, spleen, and lung.
15 In fetal tissue, the message is expressed uniformly in all tissues examined.

Table 2 Tissue distribution of human and murine β -secretase transcripts

Tissue/Organ	Size Messages Found (Kb):		Clontech Human Brain region Tissue/Organ	Human
	Human	Mouse		
Heart	2 ^a	3.5, 3.8, 5 & 7	Cerebellum	2Kb, 4Kb, 6Kb
Brain	2, 3, 4, and 7	3.5, 3.8, 5 & 7	Cerebral Cx	2Kb, 4Kb, 6Kb
Liver	2, 3, 4, and 7	3.5, 3.8, 5 & 7	Medulla	2Kb, 4Kb, 6Kb
Pancreas	2, 3, 4, and 7	nd ^d	Spinal Cord	2Kb, 4Kb, 6Kb
Placenta	2 ^a , 4 and 7 ^b	nd	Occipital Pole	2Kb, 4Kb, 6Kb
Lung	2 ^a , 4 and 7 ^b	3.5, 3.8, 5 & 7	Frontal Lobe	2Kb, 4Kb, 6Kb
Muscle	2 ^a and 7 ^b	3.5, 3.8, 5 & 7	Amygdala	2Kb, 4Kb, 6Kb
Uterus	2 ^a , 4, and 7	nd	Caudate N.	2Kb, 4Kb, 6Kb
Bladder	2 ^a , 3, 4, and 7	nd	Corpus Callosum	2Kb, 4Kb, 6Kb
Kidney	2 ^a , 3, 4, and 7	3.5, 3.8, 5 & 7	Hippocampus	2Kb, 4Kb, 6Kb
Spleen	2 ^a , 3, 4, and 7	nd		
Testis	nd	4.5Kb, 2Kb	Substantia Nigra	2Kb, 4Kb, 6Kb
Stomach	nd	5 ^a	Thalamus	2Kb, 4Kb, 6Kb
Sm. Intestine	nd	3.5, 3.8, 5 & 7		
f Brain ^c	2 ^a , 3, 4, and 7	nd		
f Liver	2 ^a , 3, 4, and 7	nd		
f Lung	2 ^a , 3, 4, and 7	nd		
f Muscle	2 ^a , 3, 4, and 7	nd		
f Heart	2 ^a , 3, 4, and 7	nd		
f Kidney	2 ^a , 3, 4, and 7	nd		
f Skin	2 ^a , 3, 4, and 7	nd		
f Sm. Intestine	2 ^a , 3, 4, and 7	nd		
Cell Line	Human	Mouse		
IMR32	2 ^a , 5 & 7			
U937	2 ^a			
THP1	2 ^a			
Jurkat	2 ^a			
HL60	none			
A293	5 & 7			
NALM6	5 & 7			
A549	5 & 7			
Hela	2, 4, 5, & 7			
PC12		2 & 5		
J774		5Kb, 2Kb		
P388D1 ccl46		5Kb (very little), 2Kb		
P19		5Kb, 2Kb		
RBL		5Kb, 2Kb		
EL4		5Kb, 2Kb		

^aby oligo 3460 probe only ^cf = fetal^bfaint ^dnd=not determined

5. Active Forms of β -secretase

a. N-terminus

The full-length open reading frame (ORF) of human β -secretase is described above, and its sequence is shown in FIG. 2A as SEQ ID NO: 2. However, as mentioned above, a further discovery of the present invention indicates that the predominant form of the active, naturally occurring molecule is truncated at the N-terminus by about 45 amino acids. That is, the protein purified from natural sources was N-terminal sequenced according to methods known in the art (Argo Bioanalytica, Morris Plains, NJ.). The N-terminus yielded the following sequence: ETDEEPEEPGRRGSFVEMVDNLRG... (SEQ ID NO: 55). This corresponds to amino acids 46-69 of the ORF-derived putative sequence. Based on this observation and others described below, the N-terminus of an active, naturally occurring, predominant human brain form of the enzyme is amino acid 46, with respect to SEQ ID NO: 2. Further processing of the purified protein provided the sequence of an internal peptide: ISFAVSACHVHDEFR (SEQ ID NO: 56), which is amino terminal to the putative transmembrane domain, as defined by the ORF. These peptides were used to validate and provide reading frame information for the isolated clones described elsewhere in this application.

In additional studies carried out in support of the present invention, N-terminal sequencing of β -secretase isolated from additional cell types revealed that the N-terminus may be amino acid numbers 46, 22, 58, or 63 with respect to the ORF sequence shown in FIG. 2A, depending on the tissue from which the protein is isolated, with the form having as its N-terminus amino acid 46 predominating in the tissues tested. That is, in experiments carried out in support of the present invention, the full-length β -secretase construct (i.e., encoding SEQ ID NO: 2) was transfected into 293T cells and COS A2 cells, using the Fugene technique described in Example 6. β -secretase was isolated from the cells by preparing a crude particulate fraction from the cell pellet, as described in Example 5, followed by extraction with buffer containing 0.2% Triton X-100. The Triton extract was diluted with pH 5.0 buffer and passed through a SP Sepharose column, essentially according to the methods described in Example 5A. This step removed the majority of contaminating proteins. After adjusting the pH to 4.5, β -secretase was further purified and concentrated on P10-P4'staD \rightarrow V Sepharose, as described in Examples 5 and 7. Fractions were analyzed for N-

terminal sequence, according to standard methods known in the art. Results are summarized in Table 3, below.

The primary N-terminal sequence of the 293T cell-derived protein was the same as that obtained from brain. In addition, minor amounts of protein starting just after the signal sequence (at Thr-22) and at the start of the aspartyl protease homology domain (Met-63) were also observed. An additional major form found in Cos A2 cells resulted from a Gly-58 cleavage.

Table 3
N-terminal Sequences and Amounts of β -secretase Forms in Various Cell Types

Source	Est. Amount (pmoles)	N-terminus (Ref.: SEQ ID NO: 2)	Sequence
Human brain	1-2	46	ETDEEPEEPGR...
Recombinant, 293T	~35	46	ETDEEPEEPGR...
	~7	22	TQHGR(LP)LR...
	~5	63	MVDNLRGKS...
Recombinant, CosA2	~4	46	ETDEEPEEPGR...
	~3	58	GSFVEMVDNL...

b. C-terminus

Further experiments carried out in support of the present invention revealed that the C-terminus of the full-length amino acid sequence presented as SEQ ID NO: 2 can also be truncated, while still retaining β -secretase activity of the molecule. More specifically, as described in more detail in Part D below, C-terminal truncated forms of the enzyme ending just before the putative transmembrane region, *i.e.* at or about 10 amino acids C terminal to amino acid 452 with respect to SEQ ID NO: 2, exhibit β -secretase activity, as evidenced by an ability to cleave APP at the appropriate cleavage site and/or ability to bind SEQ ID NO: 72.

Thus, using the reference amino acid positions provided by SEQ ID NO: 2, one form of β -secretase extends from position 46 to position 501 (β -secretase 46-501; SEQ ID NO: 43). Another form extends from position 46 to any position including and beyond position 452, (β -secretase 4-452+), with a preferred form being β -secretase 46-452 (SEQ ID NO: 58). More generally, another preferred form extends from position 1 to any position including and beyond position 452, but not including position 501. Other active forms of the β -secretase protein begin at amino acid 22, 58, or 63 and may extend to any point including and beyond the cysteine at position 420, and more preferably, including and beyond position

5 452, while still retaining enzymatic activity (i.e., β -secretase 22-452+; β -secretase 58-452+; β -secretase 63-452+). As described in Part D, below, those forms which are truncated at a
10 C-terminal position at or before about position 452, or even several amino acids thereafter, are particularly useful in crystallization studies, since they lack all or a significant portion of
5 the transmembrane region, which may interfere with protein crystallization. The recombinant protein extending from position 1 to 452 has been affinity purified using the procedures described herein.

15 C. Crystallization of β -secretase

According to a further aspect, the present invention also includes purified β -secretase
10 in crystallized form, in the absence or presence of binding substrates, such as peptide, modified peptide, or small molecule inhibitors. This section describes methods and utilities of such compositions.

1. Crystallization of the Protein

25 β -secretase purified as described above can be used as starting material to determine a
15 crystallographic structure and coordinates for the enzyme. Such structural determinations are particularly useful in defining the conformation and size of the substrate binding site. This information can be used in the design and modeling of substrate inhibitors of the enzyme. As
30 discussed herein, such inhibitors are candidate molecules for therapeutics for treatment of Alzheimer's disease and other amyloid diseases characterized by A β peptide amyloid
20 deposits.

35 The crystallographic structure of β -secretase is determined by first crystallizing the purified protein. Methods for crystallizing proteins, and particularly proteases, are now well known in the art. The practitioner is referred to Principles of Protein X-ray Crystallography
(J. Drenth, Springer Verlag, NY, 1999) for general principles of crystallography.

40 25 Additionally, kits for generating protein crystals are generally available from commercial providers, such as Hampton Research (Laguna Niguel, CA). Additional guidance can be obtained from numerous research articles that have been written in the area of crystallography
45 of protease inhibitors, especially with respect to HIV-1 and HIV-2 proteases, which are aspartic acid proteases.

30 Although any of the various forms of β -secretase described herein can be used for crystallization studies, particularly preferred forms lack the first 45 amino acids of the full
50 length sequence shown as SEQ ID NO: 2, since this appears to be the predominant form

5 which occurs naturally in human brain. It is thought that some form of post-translational
modification, possibly autocatalysis, serves to remove the first 45 amino acids in fairly rapid
order, since, to date, virtually no naturally occurring enzyme has been isolated with all of the
10 first 45 amino acids intact. In addition, it is considered preferable to remove the putative
5 transmembrane region from the molecule prior to crystallization, since this region is not
necessary for catalysis and potentially could render the molecule more difficult to crystallize.

15 Thus, a good candidate for crystallization is β -secretase 46-452 (SEQ ID NO: 58),
since this is a form of the enzyme that (a) provides the predominant naturally occurring N-
terminus, and (b) lacks the "sticky" transmembrane region, while (c) retaining β -secretase
10 activity. Alternatively, forms of the enzyme having extensions that extend part of the way
(approximately 10-15 amino acids) into the transmembrane domain may also be used. In
20 general, for determining X-ray crystallographic coordinates of the ligand binding site, any
form of the enzyme can be used that either (i) exhibits β -secretase activity, and/or (ii) binds to
a known inhibitor, such as the inhibitor ligand P10—P4'staD->V, with a binding affinity that
25 is at least 1/100 the binding affinity of β -secretase [46-501](SEQ ID NO. 43) to P10—
P4'staD->V. Therefore, a number of additional truncated forms of the enzyme can be used
in these studies. Suitability of any particular form can be assessed by contacting it with the
30 P10—P4'staD->V affinity matrix described above. Truncated forms of the enzyme that bind
to the matrix are suitable for such further analysis. Thus, in addition to 46-452, discussed
20 above, experiments in support of the present invention have revealed that a truncated form
ending in residue 419, most likely 46-419, also binds to the affinity matrix and is therefore
35 an alternate candidate protein composition for X-ray crystallographic analysis of β -secretase.
More generally, any form of the enzyme that ends before the transmembrane domain,
particularly those ending between about residue 419 and 452 are suitable in this regard.

40 25 At the N-terminus, as described above, generally the first 45 amino acids will be
removed during cellular processing. Other suitable naturally occurring or expressed forms
are listed in Table 3 above. These include, for example, a protein commencing at residue 22,
one commencing at residue 58 and one commencing at residue 63. However, analysis of the
45 entire enzyme, starting at residue 1, can also provide information about the enzyme. Other
30 forms, such as 1-420 (SEQ ID NO 60) to 1-452 (SEQ ID NO: 59), including intermediate
forms, for example 1-440, can be useful in this regard. In general, it will also be useful to
50 obtain structure on any subdomain of the active enzyme.

5 Methods for purifying the protein, including active forms, are described above. In addition, since the protein is apparently glycosylated in its naturally occurring (and mammalian-expressed recombinant) forms, it may be desirable to express the protein and
10 purify it from bacterial sources, which do not glycosylate mammalian proteins, or express it in sources, such as insect cells, that provide uniform glycosylation patterns, in order to obtain a homogeneous composition. Appropriate vectors and codon optimization procedures for accomplishing this are known in the art.

15 Following expression and purification, the protein is adjusted to a concentration of about 1-20 mg/ml. In accordance with methods that have worked for other crystallized
20 proteins, the buffer and salt concentrations present in the initial protein solution are reduced to as low a level as possible. This can be accomplished by dialyzing the sample against the starting buffer, using microdialysis techniques known in the art. Buffers and crystallization conditions will vary from protein to protein, and possibly from fragment to fragment of the active β -secretase molecule, but can be determined empirically using, for example, matrix
25 methods for determining optimal crystallization conditions. (Drentz, J., *supra*; Ducruix, A., *et al.*, eds. *Crystallization of Nucleic Acids and Proteins: A Practical Approach*, Oxford University Press, New York, 1992.)

30 Following dialysis, conditions are optimized for crystallization of the protein. Generally, methods for optimization may include making a "grid" of 1 μ l drops of the protein
35 solution, mixed with 1 μ l well solution, which is a buffer of varying pH and ionic strength. These drops are placed in individual sealed wells, typically in a "hanging drop" configuration, for example in commercially available containers (Hampton Research, Laguna Niguel, CA). Precipitation/crystallization typically occurs between 2 days and 2 weeks. Wells are checked for evidence of precipitation or crystallization, and conditions are
40 optimized to form crystals. Optimized crystals are not judged by size or morphology, but rather by the diffraction quality of crystals, which should provide better than 3 Å resolution. Typical precipitating agents include ammonium sulfate (NH_4SO_4), polyethylene glycol (PEG) and methyl pentane diol (MPD). All chemicals used should be the highest grade possible
45 (e.g., ACS) and may also be re-purified by standard methods known in the art, prior to use.

50 Exemplary buffers and precipitants forming an empirical grid for determining crystallization conditions are commercially available. For example, the "Crystal Screen" kit (Hampton Research) provides a sparse matrix method of trial conditions that is biased and selected from known crystallization conditions for macromolecules. This provides a "grid"

for quickly testing wide ranges of pH, salts, and precipitants using a very small sample (50 to 100 microliters) of macromolecule. In such studies, 1 μ l of buffer/precipitant(s) solution is added to an equal volume of dialyzed protein solution, and the mixtures are allowed to sit for at least two days to two weeks, with careful monitoring of crystallization. Chemicals can be obtained from common commercial suppliers; however, it is preferable to use purity grades suitable for crystallization studies, such as are supplied by Hampton Research (Laguna Niguel, CA). Common buffers include Citrate, TEA, CHES, Acetate, ADA and the like (to provide a range of pH optima), typically at a concentration of about 100 mM. Typical precipitants include $(\text{NH}_4)_2\text{SO}_4$, MgSO_4 , NaCl, MPD, Ethanol, polyethylene glycol of various sizes, isopropanol, KCl; and the like (Ducruix).

Various additives can be used to aid in improving the character of the crystals, including substrate analogs, ligands, or inhibitors, as discussed in Part 2, below, as well as certain additives, including, but not limited to:

- 5 % Jeffamine
- 5 % Polypropyleneglycol P400
- 5 % Polyethyleneglycol 400
- 5 % ethyleneglycol
- 5 % 2-methyl-2,4-pentanediol
- 5 % Glycerol
- 5 % Dioxane
- 5 % dimethyl sulfoxide
- 5 % n-Octanol
- 100 mM $(\text{NH}_4)_2\text{SO}_4$
- 100 mM CsCl
- 100 mM CoSO_4
- 100 mM MnCl_2
- 100 mM KCl
- 100 mM ZnSO_4
- 100 mM LiCl_2
- 100 mM MgCl_2
- 100 mM Glucose
- 100 mM 1,6-Hexanediol 100 mM Dextran sulfate
- 100 mM 6-amino caproic acid
- 100 mM 1,6 hexane diamine
- 100 mM 1,8 diamino octane
- 100 mM Spermidine
- 100 mM Spermine
- 0.17 mM n-dodecyl- β -D-maltoside NP 40
- 20 mM n-octyl- β -D-glucopyranoside

According to one discovery of the present invention, the full-length β -secretase enzyme contains at least one transmembrane domain, and its purification is aided by the use

5 of a detergent (Triton X-100). Membrane proteins can be crystallized intact, but may require specialized conditions, such as the addition of a non-ionic detergent, such as C₈G (8-alkyl-β-glucoside) or an n-alkyl-maltoside (C₈M). Selection of such a detergent is somewhat
10 empirical, but certain detergents are commonly employed. A number of membrane proteins have been successfully "salted out" by addition of high salt concentrations to the mixture.
5 PEG has also been used successfully to precipitate a number of membrane proteins (Ducruix, *et al., supra*). Alternatively, as discussed above, a C-terminal truncated form of the protein
15 that binds inhibitor but which lacks the transmembrane domain, such as β-secretase 46-452, is crystallized.

10 After crystallization conditions are determined, crystallization of a larger amount of the protein can be achieved by methods known in the art, such as vapor diffusion or
20 equilibrium dialysis. In vapor diffusion, a drop of protein solution is equilibrated against a larger reservoir of solution containing precipitant or another dehydrating agent. After sealing, the solution equilibrates to achieve supersaturating concentrations of proteins and thereby
25 induce crystallization in the drop.

Equilibrium dialysis can be used for crystallization of proteins at low ionic strength. Under these conditions, a phenomenon known as "salting in" occurs, whereby the protein
30 molecules achieve balance of electrostatic charges through interactions with other protein molecules. This method is particularly effective when the solubility of the protein is low at the lower ionic strength. Various apparatuses and methods are used, including
20 microdiffusion cells in which a dialysis membrane is attached to the bottom of a capillary tube, which may be bent at its lower portion. The final crystallization condition is achieved by slowly changing the composition of the outer solution. A variation of these methods
35 utilizes a concentration gradient equilibrium dialysis set up. Microdiffusion cells are available from commercial suppliers such as Hampton Research (Laguna Niguel, CA).
40 25

Once crystallization is achieved, crystals characterized for purity (e.g., SDS-PAGE) and biological activity. Larger crystals (>0.2 mm) are preferred to increase the resolution of
45 the X-ray diffraction, which is preferably on the order of 10-1.5 Angstroms. The selected crystals are subjected to X-ray diffraction, using a strong, monochromatic X-ray source, such
30 as a Synchrotron source or rotating anode generator, and the resulting X-ray diffraction patterns are analyzed, using methods known in the art.

50 In one application, β-secretase amino acid sequence and/or X-ray diffraction data is recorded on computer readable medium, by which is meant any medium that can be read and

5 directly accessed by a computer. These data may be used to model the enzyme, a
subdomain thereof, or a ligand thereof. Computer algorithms useful for this application are
publicly and commercially available.

10 2. Crystallization of Protein plus Inhibitor

As mentioned above, it is advantageous to co-crystallize the protein in the presence of
a binding ligand, such as inhibitor. Generally, the process for optimizing crystallization of
15 the protein is followed, with addition of greater than 1 mM concentration of the inhibitor
ligand during the precipitation phase. These crystals are also compared to crystals formed in
10 the absence of ligand, so that measurements of the ligand binding site can be made.

Alternatively, 1-2 μ l of 0.1-25 mM inhibitor compound is added to the drop containing
crystals grown in the absence of inhibitor in a process known as "soaking." Based on the
coordinates of the binding site, further inhibitor optimization is achieved. Such methods
have been used advantageously in finding new, more potent inhibitors for HIV proteases
25 (See, *e.g.*, Viswanadhan, V.N., *et al.* J. Med. Chem. 39: 705-712, 1996; Muegge, I., *et al.* J.
Med. Chem. 42: 791-804, 1999).

One inhibitor ligand which is used in these co-crystallization and soaking experiments
30 is P10—P4'staD->V (SEQ ID NO: 72), a statin peptide inhibitor described above. Methods
for making the molecule are described herein. The inhibitor is mixed with β -secretase, and
20 the mixture is subjected to the same optimization tests described above, concentrating on
those conditions worked out for the enzyme alone. Coordinates are determined and
35 comparisons are made between the free and ligand bound enzyme, according to methods well
known in the art. Further comparisons can be made by comparing the inhibitory
concentrations of the enzyme to such coordinates, such as described by Viswanadhan, *et al.*,
40 25 *supra*. Analysis of such comparisons provides guidance for design of further inhibitors,
using this method.

45 D. Biological Activity of β -secretase

1. Naturally occurring β -secretase

30 In studies carried out in support of the present invention, isolated, purified forms of β -
secretase were tested for enzymatic activity using one or more native or synthetic substrates.
50 For example, as discussed above, when β -secretase was prepared from human brain and

5 purified to homogeneity using the methods described in Example 5A, a single band was
observed by silver stain after electrophoresis of sample fractions from the anion exchange
10 chromatography (last step) on an SDS-polyacrylamide gel under reducing (+ β -
mercaptoethanol) conditions. As summarized in Table 1, above, this fraction yielded a
5 specific activity of approximately 1.5×10^9 nM/h/mg protein, where activity was measured by
hydrolysis of MBP-C125SW.

15 2. Isolated Recombinant β -secretase

Various recombinant forms of the enzyme were produced and purified from
transfected cells. Since these cells were made to overproduce the enzyme, it was found that
10 the purification scheme described with respect naturally occurring forms of the enzyme (e.g.,
Example 5A) could be shortened, with positive results. For example, as detailed in Example
20 6, 293T cells were transfected with pCEKclone 27 (FIG. 12 and FIG. 13A-E) and Cos A2
cells were transfected with pCF β A2 using "FUGENE" 6 Transfection Reagent (Roche
Molecular Biochemicals Research, Indianapolis, IN). The vector pCF was constructed from
25 the parent vector pCDNA3, commercially available from Invitrogen, by inserting SEQ ID
NO: 80 (FIG. 11A) between the HindIII and EcoRI sites. This sequence encompasses the
adenovirus major late promoter tripartite leader sequence, and a hybrid splice created from
adenovirus major late region first exon and intron and a synthetically generated IgG variable
30 region splice acceptor.

20 pCDNA3 was cut with restriction endonucleases HindIII and EcoRI, then blunted by
filling in the ends with Klenow fragment of DNA polymerase I. The cut and blunted vector
35 was gel purified, and ligated with isolated fragment from pED.GI. The pED fragment was
prepared by digesting with PvuII and SmaI, followed by gel purification of the resulting 419
base-pair fragment, which was further screened for orientation, and confirmed by sequencing.

40 25 To create the pCEK expression vector, the expression cassette from pCF was
transferred into the EBV expression vector pCEP4 (Invitrogen, Carlsbad, CA). pCEP 4 was
cut with BglII and XbaI, filled in, and the large 9.15 kb fragment containing pBR,
hygromycin, and EBV sequences) ligated to the 1.9 kb NruI to XmnI fragment of pCF
45 containing the expression cassette (CMV, TPL/MLP/IgG splice, Sp6, SVpolyA, M13
flanking region). pCF β A2 (clone A2) contains full length β -secretase in the vector pCF.
30 pCF vector replicates in COS and 293T cells. In each case, cells were pelleted and a crude
particulate fraction was prepared from the pellet. This fraction was extracted with buffer
50 containing 0.2% Triton X-100. The Triton extract was diluted with pH 5.0 buffer and passed

5 through a SP Sepharose column. After the pH was adjusted to 4.5, β -secretase activity
containing fractions were concentrated, with some additional purification on P10-
P4'(statine)D->V Sepharose, as described for the brain enzyme. Silver staining of fractions
10 revealed co-purified bands on the gel. Fractions corresponding to these bands were subjected
5 to N-terminal amino acid determination. Results from these experiments revealed some
heterogeneity of β -secretase species within the fractions. These species represent various
15 forms of the enzyme; for example, from the 293T cells, the primary N-terminus is the same as
that found in the brain, where (with respect to SEQ ID NO: 2) amino acid 46 is at the N-
terminus. Minor amounts of protein starting just after the signal sequence (at residue 23) and
10 at the start of the aspartyl protease homology domain (Met-63) were also observed. An
additional major form of protein was found in Cos A2 cells, resulting from cleavage at Gly-
20 58. These results are summarized in Table 3, above.

2. Comparison of Isolated, Naturally Occurring β -secretase with
Recombinant

15 β -secretase

As described above, naturally occurring β -secretase derived from human brain as well
as recombinant forms of the enzyme exhibit activity in cleaving APP, particularly as
30 evidenced by activity in the MBP-C125 assay. Further, key peptide sequences from the
naturally occurring form of the enzyme match portions of the deduced sequence derived from
20 cloning the enzyme. Further confirmation that the two enzymes act identically can be taken
from additional experiments in which various inhibitors were found to have very similar
35 affinities for each enzyme, as estimated by a comparison of IC_{50} values measured for each
enzyme under similar assay conditions. These inhibitors were discovered in accordance with
a further aspect of the invention, which is described below. Significantly, the inhibitors
40 25 produce near identical IC_{50} values and rank orders of potency in brain-derived and
recombinant enzyme preparations, when compared in the same assay.

In further studies, comparisons were made between the full length recombinant
enzyme having a C-terminal flag sequence "FLp501" (SEQ ID NO: 2, + SEQ ID NO: 45) and
45 a recombinant enzyme truncated at position 452 "452Stop" (SEQ ID NO: 58 or SEQ ID NO:
30 59). Both enzymes exhibited activity in cleaving β -secretase substrates such as MBP-C125,
as described above. The C-terminal truncated form of the enzyme exhibited activity in
50 cleaving the MBP-C125sw substrate as well as the P26-P4' substrate, with similar rank order

5 of potency for the various inhibitor drugs tested. In addition, the absolute IC_{50} s were comparable for the two enzymes tested with the same inhibitor. All IC_{50} s were less than 10 μ M.

10 1. Cellular β -secretase

5 Further experiments carried out in support of the present invention have revealed that the isolated β -secretase polynucleotide sequences described herein encode β -secretase or β -secretase fragments that are active in cells. This section describes experiments carried out in support of the present invention, cells were transfected with DNA encoding β -secretase alone, or were co-transfected with DNA encoding-secretase and DNA encoding wild-type APP as detailed in Example 8.

20 a. Transfection with β -secretase

In experiments carried out in support of the present invention, clones containing genes expressing the full-length polypeptide (SEQ ID NO: 2) were transfected into COS cells (Fugene and Effectene methods). Whole cell lysates were prepared and various amounts of lysate were tested for β -secretase activity according to standard methods known in the art or described in Example 4 herein. FIG. 14B shows the results of these experiments. As shown, lysates prepared from transfected cells, but not from mock- or control cells, exhibited considerable enzymatic activity in the MPB-C125sw assay, indicating "overexpression" of β -secretase by these cells.

20 b. Co-transfection of Cells with β -secretase and APP

35 In further experiments, 293T cells were co-transfected with pCEK clone 27, Figures 12 and 13 or poCK vector containing the full length β -secretase molecule (1-501; SEQ ID NO: 2) and with a plasmid containing either the wild-type or Swedish APP construct pohCK751, as described in Example 8. β -specific cleavage was analyzed by ELISA and Western analyses to confirm that the correct site of cleavage occurs.

40 Briefly, 293T cells were co-transfected with equivalent amounts of plasmids encoding β APPsw or wt and β -secretase or control β -galactosidase (β -gal) cDNA according to standard methods. β APP and β -secretase cDNAs were delivered via vectors, pohCK or pCEK, which do not replicate in 293T cells (pCEK-clone 27, FIGs. 12 and 13; pohCK751 expressing β APP 751, FIG. 21). Conditioned media and cell lysates were collected 48 hours after transfection. Western assays were carried out on conditioned media and cell lysates. ELISAs for

5 detection of A β peptide were carried out on the conditioned media to analyze various APP cleavage products.

Western Blot Results

10 It is known that β -secretase specifically cleaves at the Met-Asp in APPwt and the
5 Leu-Asp in APPsw to produce the A β peptide, starting at position 1 and releasing soluble APP (sAPP β). Immunological reagents, specifically antibody 92 and 92sw (or 192sw),
15 respectively, have been developed that specifically detect cleavage at this position in the APPwt and APPsw substrates, as described in U.S. Patent 5,721,130, incorporated herein by reference. Western blot assays were carried out on gels on which cell lysates were separated.
20 These assays were performed using methods well known in the art, using as primary antibody reagents Ab 92 or Ab92S, which are specific for the C terminus of the N-terminal fragment of APP derived from APPwt and APPsw, respectively. In addition, ELISA format assays were performed using antibodies specific to the N terminal amino acid of the C terminal fragment.

25 Monoclonal antibody 13G8 (specific for C-terminus of APP -- epitope at positions
15 675-695 of APP695) was used in a Western blot format to determine whether the transfected cells express APP. FIG.15A shows that reproducible transfection was obtained with expression levels of APP in vast excess over endogenous levels (triplicate wells are indicated as 1, 2, 3 in FIG.15A). Three forms of APP -- mature, immature and endogenous - can be
30 seen in cells transfected with APPwt or APPsw. When β -secretase was co-transfected with
20 APP, smaller C-terminal fragments appeared in triplicate well lanes from co-transfected cells (Western blot FIG. 15A, right-most set of lanes). In parallel experiments, where cells were
35 co-transfected with β -secretase and APPsw substrate, literally all of the mature APP was cleaved (right-most set of lanes labeled "1,2,3" of FIG. 15B). This suggests that there is extensive cleavage by β -secretase of the mature APP (upper band), which results in C-
40 terminal fragments of expected size in the lysate for cleavage at the β -secretase site. Co-
25 transfection with Swedish substrate also resulted in an increase in two different sized CTF fragments (indicated by star). In conjunction with the additional Western and ELISA results described below, these results are consistent with a second cleavage occurring on the APPsw
45 substrate after the initial cleavage at the β -secretase site.

30 Conditioned medium from the cells was analyzed for reactivity with the 192sw
antibody, which is specific for β -s-APPsw. Analysis using this antibody indicated a dramatic
50 increase in β -secretase cleaved soluble APP. This is observed in the gel illustrated in FIG.

5 16B by comparing the dark bands present in the "APPsw β sec" samples to the bands present in the "APPsw β gal" samples. Antibody specific for β -s-APPwt also indicates an increase in β -secretase cleaved material, as illustrated in FIG. 16A..

10 Since the antibodies used in these experiments are specific for the β -secretase cleavage site, the foregoing results show that p501 β -secretase cleaves APP at this site, and the overexpression of this recombinant clone leads to a dramatic enhancement of β -secretase processing at the correct β -secretase site in whole cells. This processing works on the wildtype APP substrate and is enhanced substantially on the Swedish APP substrate. Since approximately 20% of secreted APP in 293T cells is β -sAPP, with the increase observed below for APPsw, it is probable that almost all of the sAPP is β -sAPP. This observation was further confirmed by independent Western assays in which alpha and total sAPP were measured.

25 Monoclonal antibody 1736 is specific for the exposed α -secretase cleaved β -APP (Selkoe, et al.). When Western blots were performed using this antibody as primary antibody, a slight but reproducible decrease in α -cleaved APPwt was observed (FIG. 17A), and a dramatic decrease in α -cleaved APPsw material was also observed (note near absence of reactivity in FIG. 17B in the lanes labeled "APPsw β sec"). These results suggest that the overexpressed recombinant p501 β -secretase cleaves APPsw so efficiently or extensively that there is little or no substrate remaining for α -secretase to cleave. This further indicates that all the sAPP in APPsw β sec samples (illustrated in FIG 16B) is β -sAPP.

A β ELISA Results

Conditioned media from the recombinant cells was collected, diluted as necessary and tested for A β peptide production by ELISA on microtiter plates coated with monoclonal antibody 2G3, which is specific for recognizing the C-terminus of A β (1-40), with the detector reagent biotinylated mAb 3D6, which measures A β (x-40) (i.e., all N-terminus-truncated forms of the A β peptide). Overexpression of β -secretase with APPwt resulted in an approximately 8-fold increase in A β (x-40) production, with 1-40 representing a small percentage of the total. There was also a substantial increase in the production of A β 1-40 (FIG. 18). With APPsw there was an approximate 2-fold increase in A β (x-40). Without adhering to any particular underlying theory, it is thought that the less dramatic increase of A β (x-40) β -sec/APPsw cells in comparison to the β -sec/APPwt cells is due in part to the fact that processing of the APPsw substrate is much more efficient than that of the APPwt substrate. That is, a significant amount of APPsw is processed by endogenous β -secretase, so further increases upon transfection of β -secretase are therefore limited. These data indicate that the expression of recombinant β -secretase increases A β production and that β -secretase is rate limiting for production of A β in cells. This means that β -secretase enzymatic activity is rate limiting for production of A β in cells, and therefore provides a good therapeutic target.

IV. Utility

A. Expression Vectors and Cells Expressing β -secretase

The invention includes further cloning and expression of members of the aspartyl protease family described above, for example, by inserting polynucleotides encoding the proteins into standard expression vectors and transfecting appropriate host cells according to standard methods discussed below. Such expression vectors and cells expressing, for example, the human β -secretase enzyme described herein, have utility, for example, in producing components (purified enzyme or transfected cells) for the screening assays discussed in Part B, below. Such purified enzyme also has utility in providing starting materials for crystallization of the enzyme, as described in Section III, above. In particular, truncated form(s) of the enzyme, such as 1-452 (SEQ ID NO: 59) and 46-452 (SEQ ID NO:58), and the deglycosylated forms of the enzyme described herein are considered to have utility in this regard, as are other forms truncated partway into the transmembrane region, for example 1-460 or 46-458.

5 In accordance with the present invention, polynucleotide sequences which encode human β -secretase, splice variants, fragments of the protein, fusion proteins, or functional
10 equivalents thereof, collectively referred to herein as " β -secretase," may be used in recombinant DNA molecules that direct the expression of β -secretase in appropriate host
5 cells. Due to the inherent degeneracy of the genetic code, other nucleic acid sequences that encode substantially the same or a functionally equivalent amino acid sequence may be used
15 to clone and express β -secretase. Such variations will be readily ascertainable to persons skilled in the art.

The polynucleotide sequences of the present invention can be engineered in order to
10 alter a β -secretase coding sequence for a variety of reasons, including but not limited to, alterations that modify the cloning, processing and/or expression of the gene product. For
20 example, alterations may be introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to produce splice variants, etc. For example, it may be
25 advantageous to produce β -secretase -encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular prokaryotic or eukaryotic host (Murray, E. *et al.* (1989) *Nuc Acids Res* 17:477-508) can be selected, for example, to
30 increase the rate of β -secretase polypeptide expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced
20 from naturally occurring sequence. This may be particularly useful in producing recombinant enzyme in non-mammalian cells, such as bacterial, yeast, or insect cells. The present
35 invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation.
40 25 In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. Appropriate cloning and expression vectors for use with prokaryotic
45 and eukaryotic hosts are also described in Sambrook, *et al.*, (*supra*).

30 The present invention also relates to host cells that are genetically engineered with vectors of the invention, and the production of proteins and polypeptides of the invention by
50 recombinant techniques. Host cells are genetically engineered (i.e., transduced, transformed

5 or transfected) with the vectors of this invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral
10 particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying
5 the β -secretase gene. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art. Exemplary methods for transfection of various types of cells are provided
15 in Example 6, herein.

As described above, according to a preferred embodiment of the invention, host cells
10 can be co-transfected with an enzyme substrate, such as with APP (such as wild type or Swedish mutation form), in order to measure activity in a cell environment. Such host cells are of particular utility in the screening assays of the present invention, particularly for
20 screening for therapeutic agents that are able to traverse cell membranes.

The polynucleotides of the present invention may be included in any of a variety of
25 expression vectors suitable for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of
30 plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host. The appropriate DNA sequence may be inserted into the vector by a variety of
20 procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and related sub-cloning procedures are deemed to be within the scope of those skilled in the art.
35

The DNA sequence in the expression vector is operatively linked to an appropriate
40 transcription control sequence (promoter) to direct mRNA synthesis. Examples of such promoters include: CMV, LTR or SV40 promoter, the *E. coli* lac or trp promoter, the phage lambda PL promoter, and other promoters known to control expression of genes in
45 prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation, and a transcription terminator. The vector
30 may also include appropriate sequences for amplifying expression. In addition, the expression vectors preferably contain one or more selectable marker genes to provide a
50 phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or

5 neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

10 The vector containing the appropriate DNA sequence as described above, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells, such as yeast; insect cells such as *Drosophila* and *Spodoptera Sf9*; mammalian cells such as CHO, COS, BHK, HEK 293 or Bowes melanoma; adenoviruses; plant cells, etc. It is understood that not all cells or cell lines will be capable of producing fully functional β -secretase; for example, it is probable that human β -secretase is highly glycosylated in native form, and such glycosylation may be necessary for activity. In this event, eukaryotic host cells may be preferred. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein. The invention is not limited by the host cells employed.

15 In bacterial systems, a number of expression vectors may be selected depending upon the use intended for β -secretase. For example, when large quantities of β -secretase or fragments thereof are needed for the induction of antibodies, vectors, which direct high level expression of fusion proteins that are readily purified, may be desirable. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as Bluescript(R) (Stratagene, La Jolla, CA), in which the β -secretase coding sequence may be ligated into the vector in-frame with sequences for the amino-terminal Met and the subsequent 7 residues of beta-galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster (1989) J Biol Chem 264:5503-5509); pET vectors (Novagen, Madison WI); and the like.

20 In the yeast *Saccharomyces cerevisiae* a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH may be used. For reviews, see Ausubel *et al.* (supra) and Grant *et al.* (1987; Methods in Enzymology 153:516-544).

25 In cases where plant expression vectors are used, the expression of a sequence encoding β -secretase may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV (Brisson *et al.* (1984) Nature 310:511-514) may be used alone or in combination with the omega leader sequence from

5 TMV (Takamatsu *et al.* (1987) EMBO J 6:307-311). Alternatively, plant promoters such as
the small subunit of RUBISCO (Coruzzi *et al.* (1984) EMBO J 3:1671-1680; Broglie *et al.*
(1984) Science 224:838-843); or heat shock promoters (Winter J and Sinibaldi RM (1991)
10 Results. Probl. Cell Differ. 17:85-105) may be used. These constructs can be introduced into
5 plant cells by direct DNA transformation or pathogen-mediated transfection. For reviews of
such techniques, see Hobbs S or Murry LE (1992) in McGraw Hill Yearbook of Science and
Technology, McGraw Hill, New York, N.Y., pp 191-196; or Weissbach and Weissbach
15 (1988) Methods for Plant Molecular Biology, Academic Press, New York, N.Y., pp 421-463.

β -secretase may also be expressed in an insect system. In one such system,
10 *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express
foreign genes in *Spodoptera frugiperda* Sf9 cells or in *Trichoplusia* larvae. The β -secretase
20 coding sequence is cloned into a nonessential region of the virus, such as the polyhedrin gene,
and placed under control of the polyhedrin promoter. Successful insertion of Kv-SL coding
sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat
25 protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or
15 *Trichoplusia* larvae in which β -secretase is expressed (Smith *et al.* (1983) J Virol 46:584;
Engelhard EK *et al.* (1994) Proc Natl Acad Sci 91:3224-3227).

30 In mammalian host cells, a number of viral-based expression systems may be utilized.
In cases where an adenovirus is used as an expression vector, a β -secretase coding sequence
20 may be ligated into an adenovirus transcription/translation complex consisting of the late
promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the
35 viral genome will result in a viable virus capable of expressing the enzyme in infected host
cells (Logan and Shenk (1984) Proc Natl Acad Sci 81:3655-3659). In addition, transcription
enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to increase
40 25 expression in mammalian host cells.

Specific initiation signals may also be required for efficient translation of a β -
secretase coding sequence. These signals include the ATG initiation codon and adjacent
45 sequences. In cases where β -secretase coding sequence, its initiation codon and upstream
sequences are inserted into the appropriate expression vector, no additional translational
30 control signals may be needed. However, in cases where only coding sequence, or a portion
thereof, is inserted, exogenous transcriptional control signals including the ATG initiation
50 codon must be provided. Furthermore, the initiation codon must be in the correct reading

5 frame to ensure transcription of the entire insert. Exogenous transcriptional elements and
initiation codons can be of various origins, both natural and synthetic. The efficiency of
expression may be enhanced by the inclusion of enhancers appropriate to the cell system in
10 use (Scharf D *et al.* (1994) Results Probl Cell Differ 20:125-62; Bittner *et al.* (1987) Methods
5 in Enzymol 153:516-544).

In a further embodiment, the present invention relates to host cells containing the
above-described constructs. The host cell can be a higher eukaryotic cell, such as a
15 mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a
prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be
10 effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or
electroporation (Davis, L., Dibner, M., and Battey, I. (1986) Basic Methods in Molecular
20 Biology) or newer methods, including lipid transfection with "FUGENE" (Roche Molecular
Biochemicals, Indianapolis, IN) or "EFFECTENE" (Quiagen, Valencia, CA), or other DNA
carrier molecules. Cell-free translation systems can also be employed to produce
25 polypeptides using RNAs derived from the DNA constructs of the present invention.

A host cell strain may be chosen for its ability to modulate the expression of the
inserted sequences or to process the expressed protein in the desired fashion. Such
30 modifications of the protein include, but are not limited to, acetylation, carboxylation,
glycosylation, phosphorylation, lipidation and acylation. Post-translational processing which
20 cleaves a "prepro" form of the protein may also be important for correct insertion, folding
and/or function. For example, in the case of β -secretase, it is likely that the N-terminus of
35 SEQ ID NO: 2 is truncated, so that the protein begins at amino acid 22, 46 or 57-58 of SEQ
ID NO: 2. Different host cells such as CHO, HeLa, BHK, MDCK, 293, WI38, etc. have
specific cellular machinery and characteristic mechanisms for such post-translational
40 25 activities and may be chosen to ensure the correct modification and processing of the
introduced, foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression may
be preferred. For example, cell lines that stably express β -secretase may be transformed
45 using expression vectors which contain viral origins of replication or endogenous expression
30 elements and a selectable marker gene. Following the introduction of the vector, cells may be
allowed to grow for 1-2 days in an enriched media before they are switched to selective
50 media. The purpose of the selectable marker is to confer resistance to selection, and its
presence allows growth and recovery of cells that successfully express the introduced

5 sequences. Resistant clumps of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. For example, in experiments carried out in support of the present invention, overexpression of the "452stop" form of the enzyme has
10 been achieved.

5 Host cells transformed with a nucleotide sequence encoding β -secretase may be cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be
15 secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors
10 containing polynucleotides encoding β -secretase can be designed with signal sequences which direct secretion of β -secretase polypeptide through a prokaryotic or eukaryotic cell
20 membrane.

β -secretase may also be expressed as a recombinant protein with one or more
25 additional polypeptide domains added to facilitate protein purification. Such purification
15 facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS
30 extension/affinity purification system (Immunex Corp, Seattle, Wash.). The inclusion of a protease-cleavable polypeptide linker sequence between the purification domain and β -
20 secretase is useful to facilitate purification. One such expression vector provides for expression of a fusion protein comprising β -secretase (e.g., a soluble β -secretase fragment)
35 fused to a polyhistidine region separated by an enterokinase cleavage site. The histidine residues facilitate purification on IMLAC (immobilized metal ion affinity chromatography, as described in Porath *et al.* (1992) Protein Expression and Purification 3:263-281) while the
40 25 enterokinase cleavage site provides a means for isolating β -secretase from the fusion protein. pGEX vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are
45 soluble and can easily be purified from lysed cells by adsorption to ligand-agarose beads (e.g., glutathione-agarose in the case of GST-fusions) followed by elution in the presence of
30 free ligand.

50 Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g.,

5 temperature shift or chemical induction) and cells are cultured for an additional period. Cells
are typically harvested by centrifugation, disrupted by physical or chemical means, and the
resulting crude extract retained for further purification. Microbial cells employed in
10 expression of proteins can be disrupted by any convenient method, including freeze-thaw
5 cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods,
which are well known to those skilled in the art.

15 β -secretase can be recovered and purified from recombinant cell cultures by any of a
number of methods well known in the art, or, preferably, by the purification scheme described
herein. Protein refolding steps can be used, as necessary, in completing configuration of the
10 mature protein. Details of methods for purifying naturally occurring as well as purified forms
of β -secretase are provided in the Examples.

B. Methods of Selecting β -secretase Inhibitors

25 The present invention also includes methods for identifying molecules, such as
15 synthetic drugs, antibodies, peptides, or other molecules, which have an inhibitory effect on
the activity of β -secretase described herein, generally referred to as inhibitors, antagonists or
blockers of the enzyme. Such an assay includes the steps of providing a human β -secretase,
30 such as the β -secretase which comprises SEQ ID NO: 2, SEQ ID NO: 43, or more
particularly in reference to the present invention, an isolated protein, about 450 amino acid
20 residues in length, which includes an amino acid sequence that is at least 90% identical to
SEQ ID NO: 75 [63-423] including conservative substitutions thereof, which exhibits β -
35 secretase activity, as described herein. The β -secretase enzyme is contacted with a test
compound to determine whether it has a modulating effect on the activity of the enzyme, as
discussed below, and selecting from test compounds capable of modulating β -secretase
40 activity. In particular, inhibitory compounds (antagonists) are useful in the treatment of
25 disease conditions associated with amyloid deposition, particularly Alzheimer's disease.
Persons skilled in the art will understand that such assays may be conveniently transformed
into kits.

45 Particularly useful screening assays employ cells which express both β -secretase and
30 APP. Such cells can be made recombinantly by co-transfection of the cells with
polynucleotides encoding the proteins, as described in Section III, above, or can be made by
50 transfecting a cell which naturally contains one of the proteins with the second protein. In a

5 particular embodiment, such cells are grown up in multi-well culture dishes and are exposed
to varying concentrations of a test compound or compounds for a pre-determined period of
time, which can be determined empirically. Whole cell lysates, cultured media or cell
10 membranes are assayed for β -secretase activity. Test compounds which significantly inhibit
5 activity compared to control (as discussed below) are considered therapeutic candidates.

Isolated β -secretase, its ligand-binding, catalytic, or immunogenic fragments, or
15 oligopeptides thereof, can be used for screening therapeutic compounds in any of a variety of
drug screening techniques. The protein employed in such a test may be membrane-bound,
free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly.

20 The formation of binding complexes between β -secretase and the agent being tested can be
measured. Compounds that inhibit binding between β -secretase and its substrates, such as
APP or APP fragments, may be detected in such an assay. Preferably, enzymatic activity will
be monitored, and candidate compounds will be selected on the basis of ability to inhibit such
25 activity. More specifically, a test compound will be considered as an inhibitor of β -secretase
15 if the measured β -secretase activity is significantly lower than β -secretase activity measured
in the absence of test compound. In this context, the term "significantly lower" means that in
the presence of the test compound the enzyme displays an enzymatic activity which, when
30 compared to enzymatic activity measured in the absence of test compound, is measurably
lower, within the confidence limits of the assay method. Such measurements can be assessed
20 by a change in K_m and/or V_{max} , single assay endpoint analysis, or any other method standard
in the art. Exemplary methods for assaying β -secretase are provided in Example 4 herein.

35 For example, in studies carried out in support of the present invention, compounds
were selected based on their ability to inhibit β -secretase activity in the MBP-C125 assay.
Compounds that inhibited the enzyme activity at a concentration lower than about 50 μM
40 were selected for further screening.

The groups of compounds that are most likely candidates for inhibitor activity
comprise a further aspect of the present invention. Based on studies carried out in support of
45 the invention, it has been determined that the peptide compound described herein as P10-
P4'staD->V (SEQ ID NO: 72) is a reasonably potent inhibitor of the enzyme. Further studies
30 based on this sequence and peptidomimetics of portions of this sequence have revealed a
number of small molecule inhibitors.

5 Random libraries of peptides or other compounds can also be screened for
suitability as β -secretase inhibitors. Combinatorial libraries can be produced for many types
of compounds that can be synthesized in a step-by-step fashion. Such compounds include
10 polypeptides, beta-turn mimetics, polysaccharides, phospholipids, hormones, prostaglandins,
5 steroids, aromatic compounds, heterocyclic compounds, benzodiazepines, oligomeric N-
substituted glycines and oligocarbamates. Large combinatorial libraries of the compounds
can be constructed by the encoded synthetic libraries (ESL) method described in Affymax,
15 WO 95/12608, Affymax, WO 93/06121, Columbia University, WO 94/08051, Pharmacopeia,
WO 95/35503 and Scripps, WO 95/30642 (each of which is incorporated by reference for all
10 purposes).

20 A preferred source of test compounds for use in screening for therapeutics or
therapeutic leads is a phage display library. See, e.g., Devlin, WO 91/18980; Key, B.K., *et*
al., eds., Phage Display of Peptides and Proteins, A Laboratory Manual, Academic Press, San
25 Diego, CA, 1996. Phage display is a powerful technology that allows one to use phage
15 genetics to select and amplify peptides or proteins of desired characteristics from libraries
containing 10^6 - 10^9 different sequences. Libraries can be designed for selected variegation of
an amino acid sequence at desired positions, allowing bias of the library toward desired
30 characteristics. Libraries are designed so that peptides are expressed fused to proteins that are
displayed on the surface of the bacteriophage. The phage displaying peptides of the desired
20 characteristics are selected and can be regrown for expansion. Since the peptides are
amplified by propagation of the phage, the DNA from the selected phage can be readily
35 sequenced facilitating rapid analyses of the selected peptides.

Phage encoding peptide inhibitors can be selected by selecting for phage that bind
specifically to β -secretase protein. Libraries are generated fused to proteins such as gene II
40 25 that are expressed on the surface of the phage. The libraries can be composed of peptides of
various lengths, linear or constrained by the inclusion of two Cys amino acids, fused to the
phage protein or may also be fused to additional proteins as a scaffold. One may start with
libraries composed of random amino acids or with libraries that are biased to sequences in the
45 β APP substrate surrounding the β -secretase cleavage site or preferably, to the D \rightarrow V
30 substituted site exemplified in SEQ ID NO: 72. One may also design libraries biased toward
the peptidic inhibitors and substrates described herein or biased toward peptide sequences
50 obtained from the selection of binding phage from the initial libraries provide additional test
inhibitor compound.

5 The β -secretase is immobilized and phage specifically binding to the β -secretase selected for. Limitations, such as a requirement that the phage not bind in the presence of a known active site inhibitor of β -secretase (e.g. the inhibitors described herein), serve to further direct phage selection active site specific compounds. This can be complicated by a differential selection format. Highly purified β -secretase, derived from brain or preferably from recombinant cells can be immobilized to 96 well plastic dishes using standard techniques (reference phage book). Recombinant β -secretase, designed to be fused to a peptide that can bind (e.g. streptavidin binding motifs, His, FLAG or myc tags) to another protein immobilized (such as streptavidin or appropriate antibodies) on the plastic petri dishes can also be used. Phage are incubated with the bound β -secretase and unbound phage removed by washing. The phage are cluted and this selection is repeated until a population of phage binding to β -secretase is recovered. Binding and elution are carried out using standard techniques.

Alternatively β -secretase can be "bound" by expressing it in Cos or other mammalian cells growing on a petri dish. In this case one would select for phage binding to the β -secretase expressing cells, and select against phage that bind to the control cells, that are not expressing β -secretase.

One can also use phage display technology to select for preferred substrates of β -secretase, and incorporate the identified features of the preferred substrate peptides obtained by phage display into inhibitors.

In the case of β -secretase, knowledge of the amino acid sequence surrounding the cleavage site of APP and of the cleavage site of APPsw has provided information for purposes of setting up the phage display screening library to identify preferred substrates of β -secretase. As mentioned above, knowledge of the sequence of a particularly good peptide inhibitor, P10-P4staD->V, as described herein, provides information for setting up a "biased" library toward this sequence.

For example, the peptide substrate library containing 10^8 different sequences is fused to a protein (such as a gene III protein) expressed on the surface of the phage and a sequence that can be used for binding to streptavidin, or another protein, such as His tag and antibody to His. The phage are digested with protease, and undigested phage are removed by binding to appropriate immobilized binding protein, such as streptavidin. This selection is repeated until a population of phage encoding substrate peptide sequences is recovered. The DNA in

5 the phage is sequenced to yield the substrate sequences. These substrates are then used for further development of peptidomimetics, particularly peptidomimetics having inhibitory properties.

10 Combinatorial libraries and other compounds are initially screened for suitability by
5 determining their capacity to bind to, or preferably, to inhibit β -secretase activity in any of the assays described herein or otherwise known in the art. Compounds identified by such screens are then further analyzed for potency in such assays. Inhibitor compounds can then
15 be tested for prophylactic and therapeutic efficacy in transgenic animals predisposed to an amyloidogenic disease, such as various rodents bearing a human APP-containing transgene,
10 e.g., mice bearing a 717 mutation of APP described by Games et al., *Nature* 373: 523-527, 1995 and Wadsworth et al. (US 5,811,633, US 5,604,131, US 5,720,936), and mice bearing a Swedish mutation of APP such as described by McConlogue et al. (US 5,612,486) and Hsiao et al. (U.S. 5,877,399); Staufenberg et al., *Proc. Natl. Acad. Sci. USA* 94, 13287-13292 (1997);
20 Sturchler-Pierrat et al., *Proc. Natl. Acad. Sci. USA* 94, 13287-13292 (1997); Borchelt et al.,
25 *Neuron* 19, 939-945 (1997), all of which are incorporated herein by reference.

Compounds or agents found to be efficacious and safe in such animal models will be further tested in standard toxicological assays. Compounds showing appropriate
30 toxicological and pharmacokinetic profiles will be moved into human clinical trials for treatment of Alzheimer's disease and related diseases. The same screening approach can be
20 used on other potential agents such as peptidomimetics described above.

In general, in selecting therapeutic compounds based on the foregoing assays, it is
35 useful to determine whether the test compound has an acceptable toxicity profile, e.g., in a variety of in vitro cells and animal model(s). It may also be useful to search the tested and identified compound(s) against existing compound databases to determine whether the
40 25 compound or analogs thereof have been previously employed for pharmaceutical purposes, and if so, optimal routes of administration and dose ranges. Alternatively, routes of administration and dosage ranges can be determined empirically, using methods well known in the art (see, e.g., Benet, L.Z., et al. *Pharmacokinetics in Goodman & Gilman's The Pharmacological Basis of Therapeutics*, Ninth Edition, Hardman, J.G., et al., Eds., McGraw-
45 Hill, New York, 1966) applied to standard animal models, such as a transgenic PDAPP animal model (e.g., Games, D., et al. *Nature* 373: 523-527, 1995; Johnson-Wood, K., et al.,
30 *Proc. Natl. Acad. Sci. USA* 94: 1550-1555, 1997). To optimize compound activity and/or specificity, it may be desirable to construct a library of near-neighbor analogs to search for
50

5 analogs with greater specificity and/or activity. Methods for synthesizing near-neighbor and/or targeted compound libraries are well-known in the combinatorial library field.

C. Inhibitors and Therapeutics

10 Part B, above, describes method of screening for compounds having β -secretase inhibitory activity. To summarize, guidance is provided for specific methods of screening for potent and selective inhibitors of β -secretase enzyme. Significantly, the practitioner is directed to a specific peptide substrate/inhibitor sequences, such as P10-P4'staD->V, on
15 which drug design can be based and additional sources, such as biased phage display libraries, that should provide additional lead compounds.

20 The practitioner is also provided ample guidance for further refinement of the binding site of the enzyme, for example, by crystallizing the purified enzyme in accord with the methods provide herein. Noting the success in this area that has been enjoyed in the area of HIV protease inhibitor development, it is contemplated that such efforts will lead to further optimization of the test compounds described herein. With optimized compounds in hand, it
25 is possible to define a compound pharmacophore, and further search existing pharmacophore databases, *e.g.*, as provided by Tripos, to identify other compounds that may differ in 2-D structural formulae with the originally discovered compounds, but which share a common pharmacophore structure and activity. Test compounds are assayed in any of the inhibitor assays described herein, at various stages in development. Therefore, the present invention
30 includes β -secretase inhibitory agents discovered by any of the methods described herein, particularly the inhibitor assays and the crystallization/optimization protocols. Such inhibitory agents are therapeutic candidates for treatment of Alzheimer's disease, as well as other amyloidoses characterized by A β peptide deposition. The considerations concerning therapeutic index (toxicology), bioavailability and dosage discussed in Part B above are also
35 important to consider with respect to these therapeutic candidates.
40 25

D. Methods of Diagnosis

45 The present invention also provides methods of diagnosing individuals who carry mutations that provide enhanced β -secretase activity. For example, there are forms of familial Alzheimer's disease in which the underlying genetic disorder has yet to be
30 recognized. Members of families possessing this genetic predisposition can be monitored for alterations in the nucleotide sequence that encodes β -secretase and/or promoter regions
50

5 thereof, since it is apparent, in view of the teachings herein, that individuals who overexpress
of the enzyme or possess catalytically more efficient forms of the enzyme would be likely to
produce relatively more A β peptide. Support for this supposition is provided by the
10 observation, reported herein, that the amount of β -secretase enzyme is rate limiting for
5 production of A β in cells.

More specifically, persons suspected to have a predilection for developing for
developing or who already have the disease, as well as members of the general population,
15 may be screened by obtaining a sample of their cells, which may be blood cells or fibroblasts,
for example, and testing the samples for the presence of genetic mutations in the β -secretase
10 gene, in comparison to SEQ ID NO: 1 described herein, for example. Alternatively or in
20 addition, cells from such individuals can be tested for β -secretase activity. According to this
embodiment, a particular enzyme preparation might be tested for increased affinity and/or
Vmax with respect to a β -secretase substrate such as MBP-C125, as described herein, with
25 comparisons made to the normal range of values measured in the general population.
15 Individuals whose β -secretase activity is increased compared to normal values are susceptible
to developing Alzheimer's disease or other amyloidogenic diseases involving deposition of
A β peptide.

30 E. Therapeutic Animal Models

20 A further utility of the present invention is in creation of certain transgenic and/or
knockout animals that are also useful in the screening assays described herein. Of particular
35 use is a transgenic animal that overexpresses the β -secretase enzyme, such as by adding an
additional copy of the mouse enzyme or by adding the human enzyme. Such an animal can
be made according to methods well known in the art (e.g., Cordell, U.S. Patent 5,387,742;
40 Wadsworth et al., US 5,811,633, US 5,604,131, US 5,720,936; McConlogue et al., US
25 5,612,486; Hsiao et al., U.S. 5,877,399; and "Manipulating the Mouse Embryo, A Laboratory
Manual," B. Hogan, F. Costantini and E. Lacy, Cold Spring Harbor Press, 1986)),
45 substituting the one or more of the constructs described with respect to β -secretase, herein,
for the APP constructs described in the foregoing references, all of which are incorporated by
30 reference.

50 An overexpressing β -secretase transgenic mouse will make higher levels of A β and
s β APP from APP substrates than a mouse expressing endogenous β -secretase. This would

5 facilitate analysis of APP processing and inhibition of that processing by candidate
therapeutic agents. The enhanced production of A β peptide in mice transgenic for β -secretase
would allow acceleration of AD-like pathology seen in APP transgenic mice. This result can
10 be achieved by either crossing the β -secretase expressing mouse onto a mouse displaying
5 AD-like pathology (such as the PDAPP or Hsiao mouse) or by creating a transgenic mouse
expressing both the β -secretase and APP transgene.

15 Such transgenic animals are used to screen for β -secretase inhibitors, with the
advantage that they will test the ability of such inhibitors to gain entrance to the brain and to
effect inhibition *in vivo*.

20 10 Another animal model contemplated by the present invention is a so-called "knock-
out mouse" in which the endogenous enzyme is either permanently (as described in US Patent
Nos. 5,464,764, 5,627,059 and 5,631,153, which are incorporated by reference in their entirety)
or inducibly deleted (as described in US Patent Nos. 4,959,317, which is incorporated by
25 reference in its entirety), or which is inactivated, as described in further detail below. Such
15 mice serve as controls for β -secretase activity and/or can be crossed with APP mutant mice,
to provide validation of the pathological sequelae. Such mice can also provide a screen for
other drug targets, such as drugs specifically directed at A β deposition events.

30 β -secretase knockout mice provide a model of the potential effects of β -secretase
inhibitors *in vivo*. Comparison of the effects of β -secretase test inhibitors *in vivo* to the
35 20 phenotype of the β -secretase knockout can help guide drug development. For example, the
phenotype may or may not include pathologies seen during drug testing of β -secretase
inhibitors. If the knockout does not show pathologies seen in the drug-treated mice, one
could infer that the drug is interacting non-specifically with another target in addition to the
40 β -secretase target. Tissues from the knockout can be used to set up drug binding assays or to
25 carry out expression cloning to find the targets that are responsible for these toxic effects.
Such information can be used to design further drugs that do not interact with these
45 undesirable targets. The knockout mice will facilitate analyses of potential toxicities that are
inherent to β -secretase inhibition. Knowledge of potential toxicities will help guide the
design of design drugs or drug-delivery systems to reduce such toxicities. Inducible knockout
30 mice are particularly useful in distinguishing toxicity in an adult animal from embryonic

5 effects seen in the standard knockout. If the knockout confers fetal-lethal effects, the inducible knockout will be advantageous.

10 Methods and technology for developing knock-out mice have matured to the point that a number of commercial enterprises generate such mice on a contract basis (e.g., Lexicon
5 Genetics, Woodland TX; Cell & Molecular Technologies, Lavallete, NJ; Crysaliis, DNX Transgenic Sciences, Princeton, NJ). Methodologies are also available in the art. (See Galli-Taliadoros, L.A., *et al.*, J. Immunol. Meth. 181: 1-15, 1995). Briefly, a genomic clone of the
15 enzyme of interest is required. Where, as in the present invention, the exons encoding the regions of the protein have been defined, it is possible to achieve inactivation of the gene
10 without further knowledge of the regulatory sequences controlling transcription. Specifically, a mouse strain 129 genomic library can be screened by hybridization or PCR, using the
20 sequence information provided herein, according to methods well known in the art. (Ausubel; Sambrook) The genomic clone so selected is then subjected to restriction mapping and partial
25 exonic sequencing for confirmation of mouse homologue and to obtain information for knock-out vector construction. Appropriate regions are then sub-cloned into a "knock-out"
15 vector carrying a selectable marker, such as a vector carrying a *neo*^r cassette, which renders cells resistant to aminoglycoside antibiotics such as gentamycin. The construct is further
30 engineered for disruption of the gene of interest, such as by insertion of a sequence replacement vector, in which a selectable marker is inserted into an exon of the gene, where it
20 serves as a mutagen, disrupting the coordinated transcription of the gene. Vectors are then engineered for transfection into embryonic stem (ES) cells, and appropriate colonies are
35 isolated. Positive ES cell clones are micro-injected into isolated host blastocysts to generate chimeric animals, which are then bred and screened for germline transmission of the mutant allele.

40 25 According to a further preferred embodiment, β -secretase knock-out mice can be generated such that the mutation is inducible, such as by inserting in the knock-out mice a *lox*
region flanking the β -secretase gene region. Such mice are then crossed with mice bearing a
45 "Cre" gene under an inducible promoter, resulting in at least some off-spring bearing both the "Cre" and the *lox* constructs. When expression of "Cre" is induced, it serves to disrupt the
30 gene flanked by the *lox* constructs. Such a "Cre-lox" mouse is particularly useful, when it is suspected that the knock-out mutation may be lethal. In addition, it provides the opportunity
50 for knocking out the gene in selected tissues, such as the brain. Methods for generating Cre-

lox constructs are provided by U.S. Patent 4,959,317, incorporated herein by reference, and are made on a contractual basis by Lexicon Genetics, Woodlands, TX, among others.

The following examples illustrate, but in no way are intended to limit the present invention.

Example 1

Isolation of Coding Sequences for Human β -secretase

A. PCR Cloning

Poly A+ RNA from IMR human neuroblastoma cells was reverse transcribed using the Perkin-Elmer kit. Eight degenerate primer pools, each 8 fold degenerate, encoding the N and C terminal portions of the amino acid sequence obtained from the purified protein were designed (shown in Table 4; oligos 3407 through 3422). PCR reactions were composed of cDNA from 10 ng of RNA, 1.5 mM MgCl₂, 0.125 μ l AmpliTaq® Gold, 160 μ M each dNTP (plus 20 μ M additional from the reverse transcriptase reaction), Perkin-Elmer TAQ buffer (from AmpliTaq® Gold kit, Perkin-Elmer, Foster City, CA), in a 25 μ l reaction volume. Each of oligonucleotide primers 3407 through 3414 was used in combination with each of oligos 3415 through 3422 for a total for 64 reactions. Reactions were run on the Perkin-Elmer 7700 Sequence Detection machine under the following conditions: 10 min at 95°C, 4 cycles of, 45 °C annealing for 15 second, 72 °C extension for 45 second and 95 °C denaturation for 15 seconds followed by 35 cycles under the same conditions with the exception that the annealing temperature was raised to 55 °C . (The foregoing conditions are referred to herein as "Reaction 1 conditions.") PCR products were visualized on 4% agarose gel (Northern blots) and a prominent band of the expected size (68 bp) was seen in reactions, particularly with the primers 3515-3518. The 68 kb band was sequenced and the internal region coded for the expected amino acid sequence. This gave the exact DNA sequence for 22 bp of the internal region of this fragment.

Additional sequence was deduced from the efficiency of various primer pools of discrete sequence in generating this PCR product. Primer pools 3419 to 3422 gave very poor or no product, whereas pools 3415 to 3418 gave robust signal. The difference between these pools is a CTC (3415 to 3418) vs TTC (3419 to 3422) in the 3' most end of the pools. Since CTC primed more efficiently we can conclude that the reverse complement GAG is the correct codon. Since Met coding is unique it was concluded that the following codon is ATG. Thus the exact DNA sequence obtained is:

5 CCC.GGC.CGG.AGG.GGC.AGC.TTT.GTG.GAG.ATG.GT (SEQ ID NO: 49)
encoding the amino acid sequence P G R R G S F V E M V (SEQ ID NO: 50). This
sequence can be used to design exact oligonucleotides for 3 and 5' RACE PCR on either
10 cDNA or libraries or to design specific hybridization probes to be used to screen libraries.

5 Since the degenerate PCR product was found to be so robust, this reaction may also be
used as a diagnostic for the presence of clones containing this sequence. Pools of libraries
can be screened using this PCR product to indicate the presence of a clone in the pool. The
15 pools can be broken out to identify individual clones. Screening pools of known complexity
and or size can provide information on the abundance of this clone in a library or source and
10 can approximate the size of the full length clone or message.

20 For generation of a probe, PCR reactions using oligonucleotides 3458 and 3469 or
3458 (SEQ ID NO: 19) and 3468 (SEQ ID NO: 20) (Table 4) can be carried out using the 23
RACE product, clone 9C7E.35 (30 ng, clone 9C7E.35 was isolated from origene library, sec
Example 2), or cDNA generated from brain, using the standard PCR conditions (Perkin-
25 Elmer, rtPCR and AmpliTaq® Gold kits) with the following following: 25 µl reaction
volume 1.5 mM MgCl₂, 0.125 µl of AmpliTaq® Gold (Perkin-Elmer), initial 95° for 10 min
to activate the AmpliTaq® Gold, 36 cycles of 65° 15 sec 72° 45 sec 95° for 15 sec, followed
30 by 3 min at 72°. Product was purified on a Quiagen PCR purification kit and used as a
substrate for randompriming to generate a radiolabelled probe (Sambrook, *et al., supra*;
20 Amersham RediPrime® kit). This probe was used to isolate full length close pCEK clone 27
shown in FIGS. 12 and 13 (A-E).

Derivation of full length clone pCEK clone 27

40 A human primary neuronal cell library in the mammalian expression vector pCEK2
25 vector was generated using size selected cDNA, and pools of clones generated from different
sized inserts. The cDNA library for β-secretase screening was made with poly(A)⁺ RNA
isolated from primary human neuronal cells. The cloning vector was pCEK2 (FIG. 12).

pCEK2

45 Double-stranded cDNA inserts were synthesized using the cDNA Synthesis Kit from
50 Stratagene with some modifications. The inserts were then fractionated according to their
sizes. A total of five fractions were individually ligated with double-cut (NotI and XhoI)

pCEK2 and subsequently transformed into the E. Coli strain XL-10 Gold which is designed to accept very large plasmids.

The fractions of transformed E. Coli were plated on Terrific Broth agar plates containing ampicillin and let grown for 18 hours. Each fraction yielded about 200,000 colonies to give a total of one million colonies. The colonies were then scraped from the plates and plasmids isolated from them in pools of approximately 70,000 clones/pool. 70,000 clones from each pool of the library was screened for the presence of the putative β -secretase gene using the diagnostic PCR reaction (degenerate primers 3411 and 3417 shown above).

Clones from the 1.5 kb pool were screened using a radiolabeled probe generated from a 390 b.p. PCR product generated from clone 9C7E.35. For generation of a probe, PCR product was generated using 3458 and 3468 as primers and clone 9C7E.35 (30 ng) as substrate.

PCR product was used as a substrate for random priming to generate a radiolabeled probe. 180,000 clones from the 1.5 kb pool (70,000 original clones in this pool), were screened by hybridization with the PCR probe and 9 positive clones identified. Four of these clones were isolated and by restriction mapping these appear to encode two independent clones of 4 to 5 kb (clone 27) and 6 to 7 kb (clone 53) length. Sequencing of clone 27 verified that it contains a coding region of 1.5 kb. FIG. 13 (A-E) shows the sequence of pCEK clone27 (clone 27).

Table 4

SEQ ID NO.	Pool No.	Nucleotide Sequence (Degenerate substitutions are shown in parentheses)
3	3407	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAG.GAG.CC
4	3408	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAA.GAG.CC
5	3409	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAA.GAA.CC
6	3410	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAG.GAA.CC
7	3411	AGA.GAC.GA(GA).GA(GA).CC(CG).GAG.GAG.CC
8	3412	AGA.GAC.GA(GA).GA(GA).CC(CG).GAA.GAG.CC
9	3413	AGA.GAC.GA(GA).GA(GA).CC(CG).GAA.GAA.CC
10	3414	AGA.GAC.GA(GA).GA(GA).CC(CG).GAG.GAA.CC

11	3415	CG.TCA.CAG.(GA)TT.(GA)TC.AAC.CAT.CTC
12	3416	CG.TCA.CAG.(GA)TT.(GA)TC.TAC.CAT.CTC
13	3417	CG.TCA.CAG.(GA)TT.(GA)TC.CAC.CAT.CTC
14	3418	CG.TCA.CAG.(GA)TT.(GA)TC.GAC.CAT.CTC
15	3419	CG.TCA.CAG.(GA)TT.(GA)TC.AAC.CAT.TTC
16	3420	CG.TCA.CAG.(GA)TT.(GA)TC.TAC.CAT.TTC
17	3421	CG.TCA.CAG.(GA)TT.(GA)TC.CAC.CAT.TTC
18	3422	CG.TCA.CAG.(GA)TT.(GA)TC.GAC.CAT.TTC
19	3458	GAG GGG CAG CTT TGT GGA GA
20	3468	CAG.CAT.AGG.CCA.GCC.CCA.GGA.TGC.CT
21	3469	GTG.ATG.GCA.GCA.ATG.TTG.GCA.CGC

Example 2

Screening of human fetal brain cDNA library

The Origene human fetal brain Rapid-Screen™ cDNA Library Panel is provided as a 96-well format array consisting of 5000 clones (plasmid DNA) per well from a human fetal brain library. Subplates are available for each well consisting of 96 wells of 50 clones each in *E. coli*. This is an oligo-dT primed library, size-selected and unidirectionally inserted into the vector pCMV-XL3.

94 wells from the master plate were screened using PCR. The Reaction 1 Conditions described in Example 1, above, were followed, using only primers 3407 and 3416 with 30ng of plasmid DNA from each well. Two pools showed the positive 70bp band. The same primers and conditions were used to screen 1µl *E. coli* from each well of one of the subplates. *E. coli* from the single positive well was then plated onto LB/amp plates and single colonies screened using the same PCR conditions. The positive clone, about 1Kb in size, was labeled 9C7E.35. It contained the original peptide sequence as well as 5' sequence that included a methionine. The 3' sequence did not contain a stop codon, suggesting that this was not a full-length clone, consistent with Northern blot data.

5

Example 3

PCR Cloning Methods

10

3'RACE was used in experiments carried out in support of the present invention to elucidate the polynucleotide encoding human β -secretase. Methods and conditions appropriate for replicating the experiments described herein and/or determining polynucleotide sequences encoding additional members of the novel family of aspartyl proteases described herein may be found, for example, in White, B.A., ed., PCR Cloning Protocols; Humana Press, Totowa, NJ, 1997, or Ausubel, *supra*, both of which are incorporated herein by reference.

15

20

RT-PCR

25

For reverse transcription polymerase chain reaction (RT-PCR), two partially degenerate primer sets used for RT-PCR amplification of a cDNA fragment encoding this peptide. Primer set 1 consisted of DNA's #3427-3434, the sequences of which are shown in Table 5, below. Matrix RT-PCR using combinations of primers from this set with cDNA reverse transcribed from primary human neuronal cultures as template yielded the predicted 54 bp cDNA product with primers #3428 + 3433. All RT-PCR reactions employed 10-50 ng input poly-A+ RNA equivalents per reaction, and were carried out for 35 cycles employing step cycle conditions with a 95°C denaturation for 1 minute, 50°C annealing for 30 sec, and a 72°C extension for 30 sec.

30

35

The degeneracy of primers #3428 + 3433 was further broken down, resulting in primer set 2, comprising DNAs #3448-3455 (Table 5). Matrix RT-PCR was repeated using primer set 2, and cDNA reverse transcribed from poly-A+ RNA from IMR-32 human neuroblastoma cells (American Type Culture Collection, Manassas, VA), as well as primary human neuronal cultures, as template for amplification. Primers #3450 and 3454 from set 2 most efficiently amplified a cDNA fragment of the predicted size (72 bp), although primers 3450+3453, and 3450+3455 also amplified the same product, albeit at lower efficiency. A 72 bp PCR product was obtained by amplification of cDNA from IMR-32 cells and primary human neuronal cultures with primers 3450 and 3454.

40

45

5' and 3' RACE-PCR

50

Internal primers matching the upper (coding) strand for 3' Rapid Amplification of 5' Ends (RACE) PCR, and lower (non-coding) strand for 5' RACE PCR were designed and made according to methods known in the art (*e.g.*, Frohman, M. A., M. K. Dush and G. R.

55

5 Martin (1988). "Rapid production of full-length cDNAs from rare transcripts: amplification
using a single gene specific oligo-nucleotide primer." Proc. Natl. Acad. Sci. U.S.A. 85(23):
8998-9002.) The DNA primers used for this experiment (#3459 & #3460) are presented in
10 Table 4. These primers can be utilized in standard RACE-PCR methodology employing
5 commercially available templates (e.g. Marathon Ready cDNA®, Clontech Labs), or custom
tailored cDNA templates prepared from RNAs of interest as described by Frohman et al.
(*ibid.*).

15 In experiments carried out in support of the present invention, a variation of RACE
was employed to exploit an IMR-32 cDNA library cloned in the retrovirus expression vector
10 pLPCXlox, a derivative of pLNCX. As the vector junctions provide unique anchor sequences
abutting the cDNA inserts in this library, they serve the purpose of 5' and 3' anchor primers
20 in RACE methodology. The sequences of the specific 5' and 3' anchor primers we employed
to amplify β -secretase cDNA clones from the library, primers #3475 and #3476, are derived
from the DNA sequence of the vector provided by Clontech Labs, Inc., and are shown in
25 Table 3.

15 Primers #3459 and #3476 were used for 3' RACE amplification of downstream
sequences from our IMR-32 cDNA library in the vector pLPCXlox. The library had
30 previously been sub-divided into 100 pools of 5,000 clones per pool, and plasmid DNA was
isolated from each pool. A survey of the 100 pools with the primers identified as diagnostic
20 for presence of the β -secretase clone, according to methods described in Example 1, above,
provided individual pools from the library for RACE-PCR. 100 ng template plasmid from
35 pool 23 was used for PCR amplification with primers 3459+3476. Amplification was carried
out for 40 cycles using *ampli-Taq Gold*®, under the following conditions: denaturation at
95°C for 1 min, annealing at 65°C for 45 sec., and extension at 72°C for 2 min. Reaction
40 products were fractionated by agarose gel chromatography, according to methods known in
the art (Ausubel; Sambrook).

45 An approximately 1.8 Kb PCR fragment was revealed by agarose gel fractionation of
the reaction products. The PCR product was purified from the gel and subjected to DNA
sequence analysis using primer #3459. The resulting sequence, designated 23A, and the
30 predicted amino acid sequence deduced from the DNA sequence are shown in FIG. 5. Six
of the first seven deduced amino-acids from one of the reading frames of 23A were an exact
50 match with the last 7 amino-acids of the N-terminal sequence determined from the purified

5

protein, purified and sequenced in further experiments carried out in support of the present invention, from natural sources.

10

15

20

25

30

35

40

45

50

65

55

Table 5

SEQ ID NO.	DNA #	NUCLEOTIDE SEQUENCE	COMMENTS
22	3427	GAY GAR GAG CCN GAG GA	
23	3428	GAY GAR GAG CCN GAa GA	
24	3429	GAY GAR GAa CCN GAg GA	
25	3430	GAY GAR GAa CCN GAa GA	
26	3431	RTT RTC NAC CAT TTC	
27	3432	RTT RTC NAC CAT cTC	
28	3433	TCN ACC ATY TCN ACA AA	
29	3434	TCN ACC ATY TCN ACG AA	
30	3448	ata <u>ttc tag a</u> GAY GAR GAg CCa GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 1 of 4
31	3449	ata <u>ttc tag a</u> GAY GAR GAg CCg GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 2 of 4
32	3450	ata <u>ttc tag a</u> GAY GAR GAg CCc GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 3 of 4
33	3451	ata <u>ttc tag a</u> GAY GAR GAg CCl GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 4 of 4
34	3452	aca <u>cga att c</u> TT RTC NAC CAT YTC aAC AAA	breakdown of 3433, 1 of 4; tm = 50
35	3453	aca <u>cga att c</u> TT RTC NAC CAT YTC gAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 2 of 4; tm = 50
36	3454	aca <u>cga att c</u> TT RTC NAC CAT YTC cAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 3 of 4; tm = 50
37	3455	aca <u>cga att c</u> TT RTC NAC CAT YTC iAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 4 of 4; tm = 50
38	3459	aa gaG CCC GGC CGG AGG GGC A	5' upper strand primer for 3' race encodes eEPGRRG
39	3460	aaa GCT GCC CCT CCG GCC GGG	3' lower strand primer for 5' RACE
40	3475	AGC TCG TTT AGT GAA CCG TCA GAT CG	pLNCX 5' primer
41	3476	ACC TAC AGG TGG GGT CTT TCA TTC CC	pLNCX, 3' primer

5

Example 4 **β -secretase Inhibitor Assays**

10

Assays for measuring β -secretase activity are well known in the art. Particularly useful assays, summarized below, are detailed in allowed U.S. Patent 5,744,346, incorporated herein by reference.

A. Preparation of MBP-C125sw

15

1. Preparation of cells

20

25

30

35

40

45

50

55

Two 250 ml cell culture flasks containing 50 ml LBamp100 per flask were seeded with one colony per flask of E. coli pMAL-C125SW cl. 2 (E. coli expressing MBP-C125sw fusion protein). Cells were allowed to grow overnight at 37°C. Aliquots (25 ml) were seeded in 500 ml per flask of LBamp100 in 2 liter flasks, which were then allowed to grow at 30°. Optical densities were measured at 600 nm (OD₆₀₀) vs LB broth; 1.5 ml 100mM IPTG was added when the OD was ~0.5. At this point, a pre-incubation aliquot was removed for SDS-PAGE ("I"). Of this aliquot, 0.5 ml was centrifuged for 1 min in a Beckman microfuge, and the resulting pellet was dissolved in 0.5 ml 1 x LSB. The cells were incubated/induced for 5-6 hours at 30 C, after which a post-incubation aliquot ("I") was removed. Cells were then centrifuged at 9,000 rpm in a KA9.1 rotor for 10 min at 4° C. Pellets were retained and stored at -20 C.

2. Extraction of bacterial cell pellets

Frozen cell pellets were resuspended in 50 ml 0.2 M NaCl, 50mM Tris, pH 7.5, then sonicated in rosette vessal for 5 x 20 sec bursts, with 1min rests between bursts. The extract was centrifuged at 16,500 rpm in a KA18.5 rotor 30 min (39,000 x g). Using pipette as a pestle, the sonicated pellet was suspended in 50 ml urea extraction buffer (7.6 M urea, 50 mM Tris pH 7.5, 1 mM EDTA, 0.5% TX-100). The total volume was about 25 ml per flask. The suspension was then sonicated 6x 20 sec, with 1 min rests between bursts. The suspension was then centrifuged again at 16,500 rpm 30 min in the KA18.5 rotor. The resulting supernatant was added to 1.5 L of buffer consisting of 0.2 M NaCl 50 mM Tris buffer, pH 7.5, with 1% Triton X-100 (0.2M NaCl-Tris-1%Tx), and was stirred gently at 4 degrees C for 1 hour, followed by centrifugation at 9,000 rpm in KA9.1 for 30 min at 4°C. The supernatant was loaded onto a column of washed amylose (100 ml of 50% slurry; New England BioLabs). The column was washed with 0.2 M NaCl-Tris-1%TX to baseline (+10 column volumes), then with 2 column volumes 0.2M NaCl-Tris-1% reduced Triton X-100.

5 The protein was then eluted with 10 mM maltose in the same buffer. An equal volume of 6
M guanidine HCl/0.5% TX-100 was added to each fraction. Peak fractions were pooled and
diluted to a final concentration of about 2 mg/ml. The fractions were stored at -40 degrees C,
10 before dilution (20-fold, to 0.1 mg/ml in 0.15% Triton X-100). Diluted aliquots were also
5 stored at -40 C.

B. Antibody-based Assays

15 The assays described in this section are based on the ability of certain antibodies,
hereinafter "cleavage-site antibodies," to distinguish cleavage of APP by β -secretase, based
on the unique cleavage site and consequent exposure of a specific C-terminus formed by the
10 cleavage. The recognized sequence is a sequence of usually about 3-5 residues is
20 immediately amino terminal of the β amyloid peptide (β AP) produced by β -secretase
cleavage of β -APP, such as Val-Lys-Met in wild-type or Val-Asn-Leu- in the Swedish double
mutation variant form of APP. Recombinantly-expressed proteins, described below, were
used as substrates for β -secretase.

25 MBP-C125 Assay: MBP-C125 substrates were expressed in *E. coli* as a fusion
15 protein of the last 125 amino acids of APP fused to the carboxy-terminal end of maltose-
binding protein (MBP), using commercially available vectors from New England Biolabs.
30 The β -cleavage site was thus 26 amino acids downstream of the start of the C-125 region.
This latter site is recognized by monoclonal antibody SW192.

20 Recombinant proteins were generated with both the wild-type APP sequence (MBP-
C125 wt) at the cleavage site (..Val-Lys-Met-Asp-Ala..) or the "Swedish" double mutation
35 (MBP-C125 sw) (..Val-Asn-Leu-Asp-Ala..). As shown schematically in FIG. 19A, cleavage
of the intact MBP-fusion protein results in the generation of a truncated amino-terminal
fragment, with the new SW-192 Ab-positive epitope uncovered at the carboxy terminus. This
40 25 amino-terminal fragment can be recognized on Western blots with the same Ab, or,
quantitatively, using an anti-MBP capture-biotinylated SW-192 reporter sandwich format, as
shown in FIG. 19A. Anti-MBP polyclonal antibodies were raised in rabbits (Josman Labs,
45 Berkeley) by immunization with purified recombinantly expressed MBP (New England
Biolabs). Antisera were affinity purified on a column of immobilized MBP. MBP-C125 SW
30 and WT substrates were expressed in *E. coli*, then purified as described above.

50 Microtiter 96-well plates were coated with purified anti-MBP antibody (at a
concentration of 5-10 μ g/ml), followed by blocking with 2.5 g/liter human serum albumin in

5 1 g/liter sodium phosphate monobasic, 10.8 g/liter sodium phosphate dibasic, 25 g/liter
sucrose, 0.5 g/liter sodium azide, pH 7.4. Appropriately diluted β -secretase enzyme (5 μ l)
was mixed with 2.5 μ l of 2.2 μ M MBP-C125sw substrate stock, in a 50 μ l reaction mixture
10 with a final buffer concentration of 20 mM acetate buffer, pH 4.8, 0.06% Triton X-100, in
5 individual wells of a 96-well microtiter plate, and incubated for 1 hour at 37 degrees C.
Samples were then diluted 5-fold with Specimen Diluent (0.2 g/l sodium phosphate
15 monobasic, 2.15 g/l sodium phosphate dibasic, 0.5 g/l sodium azide, 8.5 g/l sodium chloride,
0.05% Triton X-405, 6 g/l BSA), further diluted 5-10 fold into Specimen Diluent on anti-
MBP coated plates, and incubated for 2 hours at room temperature. Following incubations
20 with samples or antibodies, plates were washed at least four times in TTBS (0.15 M NaCl, 50
mM Tris, pH 7.5, 0.05% Tween-20). Biotinylated SW192 antibodies were used as the
reporter. SW192 polyclonal antibodies were biotinylated using NHS-biotin (Pierce),
following the manufacturer's instruction. Usually, the biotinylated antibodies were used at
25 about 240 ng/ml, the exact concentration varying with the lot of antibodies used. Following
15 incubation of the plates with the reporter, the ELISA was developed using streptavidin-
labeled alkaline phosphatase (Boehringer-Mannheim) and 4-methyl-umbelliferyl phosphate as
fluorescent substrate. Plates were read in a Cytofluor 2350 Fluorescent Measurement
30 System. Recombinantly generated MBP-26SW (product analog) was used as a standard to
generate a standard curve, which allowed the conversion of fluorescent units into amount of
20 product generated.

35 This assay protocol was used to screen for inhibitor structures, using "libraries" of
compounds assembled onto 96-well microtiter plates. Compounds were added, in a final
concentration of 20 μ g/ml in 2% DMSO, in the assay format described above, and the extent
of product generated compared with control (2% DMSO only) β -secretase incubations, to
40 25 calculate "% inhibition." "Hits" were defined as compounds which result in >35% inhibition
of enzyme activity at test concentration. This assay can also be used to provide IC_{50} values
for inhibitors, by varying the concentration of test compound over a range to calculate from a
dose-response curve the concentration required to inhibit the activity of the enzyme by 50%.

45 Generally, inhibition is considered significant as compared to control activity in this
30 assay if it results in activity that is at least 1 standard deviation, and preferably 2 standard
deviations lower than a mean activity value determined over a range of samples. In addition,

5 a reduction of activity that is greater than about 25%, and preferably greater than about 35% of control activity may also be considered significant.

Using the foregoing assay system, 24 "hits" were identified (>30% inhibition at 50
10 μ M concentration) from the first 6336 compounds tested (0.4% hit rate). Of these 12
5 compounds had IC_{50} s less than 50 μ M, including re-screening in the P26-P4'sw assay, below.

P26-P4'sw assay. The P26-P4'sw substrate is a biotin-linked peptide of the sequence
15 (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNLDAEF (SEQ ID NO: 63). The P26-P1
standard has the sequence (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNL (SEQ ID
NO: 64), where the N-terminal "CGG" serves as a linker between biotin and the substrate in
10 both cases. Peptides were prepared by Anaspec, Inc. (San Jose, CA) using solid phase
20 synthesis with boc-amino acids. Biotin was coupled to the terminal cysteine sulfhydryl by
Anaspec, Inc. after synthesis of the peptide, using EZ-link Iodoacetyl-LC-Biotin (Pierce).
Peptides are stored as 0.8-1.0 mM stocks in 5 mM Tris, with the pH adjusted to around
neutral (pH 6.5-7.5) with sodium hydroxide.

25 15 For the enzyme assay, the substrate concentration can vary from 0 – 200 μ M.
Specifically for testing compounds for inhibitory activity, substrate concentration is 1.0 μ M.
Compounds to be tested were added in DMSO, with a final DMSO concentration of 5%; in
30 such experiments, the controls also receive 5% DMSO. Concentration of enzyme was varied,
to give product concentrations within the linear range of the ELISA assay (125 – 2000 pM,
20 after dilution). These components were incubated in 20 mM sodium acetate, pH 4.5, 0.06%
Triton X-100, at 37 °C for 1 to 3 hours. Samples were diluted 5-fold in specimen diluent
35 (145.4 mM sodium chloride, 9.51 mM sodium phosphate, 7.7 mM sodium azide, 0.05%
Triton X-405, 6 gm/liter bovine serum albumin, pH 7.4) to quench the reaction, then diluted
further for the ELISA as needed. For the ELISA, Costar High Binding 96-well assay plates
40 25 (Corning, Inc., Corning, NY) were coated with SW 192 monoclonal antibody from clone
16A7, or a clone of similar affinity. Biotin-P26-P4' standards were diluted in specimen
diluent to a final concentration of 0 to 2 nM. Diluted samples and standards (100 μ l) are
45 incubated on the SW192 plates at 4 °C for 24 hours. The plates are washed 4 times in TTBS
buffer (150 mM sodium chloride, 25 mM Tris, 0.05 % Tween 20, pH 7.5), then incubated
30 with 0.1 ml/well of streptavidin – alkaline phosphatase (Roche Molecular Biochemicals,
Indianapolis, IN) diluted 1:3000 in specimen diluent. After incubating for one hour at room
50 temperature, the plate was washed 4 times in TTBS, as described in the previous section, and

incubated with fluorescent substrate solution A (31.2 gm/liter 2-amino-2-methyl-1-propanol, 30 mg/liter, adjusted to pH 9.5 with HCl). Fluorescent values were read after 30 minutes.

C. Assays using Synthetic Oligopeptide Substrates

This assay format is particularly useful for measuring activity of partially purified β -secretase preparations. Synthetic oligopeptides are prepared which incorporate the known cleavage site of β -secretase, and optional detectable tags, such as fluorescent or chromogenic moieties. Examples of such peptides, as well as their production and detection methods are described in allowed U.S. Patent 5,942,400, herein incorporated by reference. Cleavage products can be detected using high performance liquid chromatography, or fluorescent or chromogenic detection methods appropriate to the peptide to be detected, according to methods well known in the art. By way of example, one such peptide has the sequence SEVNL DAEF (SEQ ID NO: 52), and the cleavage site is between residues 5 and 6. Another preferred substrate has the sequence ADRGLTTRPGSGLTNIKTEEISEVNLDAE F (SEQ ID NO: 53), and the cleavage site is between residues 26 and 27.

D. β -secretase Assays of Crude Cell or Tissue Extracts

Cells or tissues were extracted in extraction buffer (20 mM HEPES, pH 7.5, 2 mM EDTA, 0.2% Triton X-100, 1 mM PMSF, 20 μ g/ml pepstatin, 10 μ g/ml E-64). The volume of extraction buffer will vary between samples, but should be at least 200 μ l per 10^6 cells. Cells can be suspended by trituration with a micropipette, while tissue may require homogenization. The suspended samples were incubated for 30 minutes on ice. If necessary to allow pipetting, unsolubilized material was removed by centrifugation at 4 degrees C, 16,000 x g (14,000 rpm in a Beckman microfuge) for 30 minutes. The supernate was assayed by dilution into the final assay solution. The dilution of extract will vary, but should be sufficient so that the protein concentration in the assay is not greater than 60 μ g/ml. The assay reaction also contained 20 mM sodium acetate, pH 4.8, and 0.06% Triton X-100 (including Triton contributed by the extract and substrate), and 220 – 110 nM MBP-C125 (a 1:10 or 1:20 dilution of the 0.1 mg/ml stock described in the protocol for substrate preparation). Reactions were incubated for 1 – 3 hours at 37degrees C before quenching with at least 5 – fold dilution in specimen diluent and assaying using the standard protocol.

5

Example 5
Purification of β -secretase

10

A. Purification of Naturally Occurring β -secretase

15

5 Human 293 cells were obtained and processed as described in U.S. Patent 5,744,346, incorporated herein by reference. (293 cells are available from the American Type Culture Collection, Manassas, VA). Frozen tissue (293 cell paste or human brain) was cut into pieces and combined with five volumes of homogenization buffer (20 mM Hepes, pH 7.5, 0.25 M sucrose, 2 mM EDTA). The suspension was homogenized using a blender and centrifuged at 10 16,000 x g for 30 min at 4°C. The supernatants were discarded and the pellets were suspended in extraction buffer (20 mM MES, pH 6.0, 0.5% Triton X-100, 150 mM NaCl, 2 mM EDTA, 5 μ g/ml leupeptin, 5 μ g/ml E64, 1 μ g/ml pepstatin, 0.2 mM PMSF) at the original volume. After vortex-mixing, the extraction was completed by agitating the tubes at 20 4°C for a period of one hour. The mixtures were centrifuged as above at 16,000 x g, and the supernatants were pooled. The pH of the extract was adjusted to 7.5 by adding ~1% (v/v) of 1 M Tris base (not neutralized).

25

30

The neutralized extract was loaded onto a wheat germ agglutinin-agarose (WGA-agarose) column pre-equilibrated with 10 column volumes of 20 mM Tris, pH 7.5, 0.5% Triton X-100, 150 mM NaCl, 2 mM EDTA, at 4°C. One milliliter of the agarose resin was 20 used for every 1 g of original tissue used. The WGA-column was washed with 1 column volume of the equilibration buffer, then 10 volumes of 20 mM Tris, pH 7.5, 100 mM NaCl, 2 mM NaCl, 2 mM EDTA, 0.2% Triton X-100 and then eluted as follows. Three-quarter 35 column volumes of 10% chitin hydrolysate in 20 mM Tris, pH 7.5, 0.5%, 150 mM NaCl, 0.5% Triton X-100, 2 mM EDTA were passed through the column after which the flow was 25 stopped for fifteen minutes. An additional five column volumes of 10% chitin hydrolysate solution were then used to elute the column. All of the above eluates were combined (pooled WGA-eluate).

40

45

The pooled WGA-eluate was diluted 1:4 with 20 mM NaOAc, pH 5.0, 0.5% Triton X-100, 2 mM EDTA. The pH of the diluted solution was adjusted to 5.0 by adding a few 30 drops of glacial acetic acid while monitoring the pH. This "SP load" was passed through a 5-ml Pharmacia HiTrap SP-column equilibrated with 20 mM NaOAc, pH 5.0, 0.5% Triton X-100, 2 mM EDTA, at 4 ml/min at 4°C.

50

55

The foregoing methods provided peak activity having a specific activity of greater than 253 nM product/ml/h/ μ g protein in the MBP-C125-SW assay, where specific activity is determined as described below, with about 1500-fold purification of the protein. Specific activity of the purified β -secretase was measured as follows. MBP C125-SW substrate was combined at approximately 220 nM in 20 mM sodium acetate, pH 4.8, with 0.06% Triton X-100. The amount of product generated was measured by the β -secretase assay, also described below. Specific activity was then calculated as:

$$\text{Specific Activity} = \frac{(\text{Product conc. nM})(\text{Dilution factor})}{(\text{Enzyme sol. vol})(\text{Incub. time h})(\text{Enzyme conc. mg/vol})}$$

The Specific Activity is thus expressed as pmoles of product produced per μ g of β -secretase per hour. Further purification of human brain enzyme was achieved by loading the SP flow through fraction on to the P10-P4'sta D->V affinity column, according to the general methods described below. Results of this purification step are summarized in Table 1, above.

B. Purification of β -secretase from Recombinant Cells

Recombinant cells produced by the methods described herein generally were made to over-express the enzyme; that is, they produced dramatically more enzyme per cell than is found to be endogenously produced by the cells or by most tissues. It was found that some of the steps described above could be omitted from the preparation of purified enzyme under these circumstances, with the result that even higher levels of purification were achieved.

CosA2 or 293 T cells transfected with β -secretase gene construct (see Example 6) were pelleted, frozen and stored at -80 degrees until use. The cell pellet was resuspended by homogenizing for 30 seconds using a handheld homogenizer (0.5 ml/pellet of approximately 10^6 cells in extraction buffer consisting of 20 mM TRIS buffer, pH 7.5, 2 mM EDTA, 0.2% Triton X-100, plus protease inhibitors: 5 μ g/ml E-64, 10 μ g/ml pepstatin, 1 mM PMSF), centrifuged at maximum speed in a microfuge (40 minutes at 4 degrees C). Pellets were suspended in original volume of extraction buffer, then stirred at 1 hour at 4 degrees C with rotation, and centrifuged again in a microfuge at maximum speed for 40 minutes. The resulting supernatant was saved as the "extract." The extract was then diluted with 20 mM sodium acetate, pH 5.0, 2 mM EDTA and 0.2% Triton X-100 (SP buffer A), and 5M NaCl was added to a final concentration of 60 mM NaCl. The pH of the solution was then adjusted

5 to pH 5.0 with glacial acetic acid diluted 1: 10 in water. Aliquots were saved ("SP load"). The
SP load was passed through a 1 ml SP HiTrap column which was pre-washed with 5 ml SP
buffer A, 5 ml SP buffer B (SP buffer A with 1 M NaCl) and 10 ml SP buffer A. An
10 additional 2 ml of 5% SP buffer B was passed through the column to displace any remaining
5 sample from the column. The pH of the SP flow-through was adjusted to pH 4.5 with 10X
diluted acetic acid. This flow-through was then applied to a P10-P4'staD->V-Sepharose
Affinity column, as described below. The column (250 μ l bed size) was pre-equilibrated
15 with at least 20 column volumes of equilibration buffer (25 mM NaCl, 0.2% Triton X-100,
0.1 mM EDTA, 25 mM sodium acetate, pH 4.5), then loaded with the diluted supernatant.
10 After loading, subsequent steps were carried out at room temperature. The column was
washed with washing buffer (125 mM NaCl, 0.2% Triton X-100, 25 mM sodium acetate, pH
20 4.5) before addition of 0.6 column bed volumes of borate elution buffer (200 mM NaCl, 0.2%
reduced Triton X-100, 40 mM sodium borate, pH 9.5). The column was then capped, and an
additional 0.2 ml elution buffer was added. The column was allowed to stand for 30 minutes.
25 15 Two bed volumes elution buffer were added, and column fractions (250 μ l) were collected.
The protein peak eluted in two fractions. 0.5 ml of 10 mg/ml pepstatin was added per
milliliter of collected fractions.

30 Cell extracts made from cells transfected with full length clone 27 (encoding SEQ ID
NO: 2; 1-501), 419stop (SEQ ID NO: 57) and 452stop (SEQ ID NO: 59) were detected by
20 Western blot analysis using antibody 264A (polyclonal antibody directed to amino acids 46-
67 of β -secretase with reference to SEQ ID NO: 2).

Example 6

Preparation of Heterologous Cells Expressing Recombinant β -secretase

40 25 Two separate clones (pCEKclone27 and pCEKclone53) were transfected into 293T or
COS(A2) cells using Fugene and Effectene methods known in the art. 293T cells were
obtained from Edge Biosystems (Gaithersburg, MD). They are KEK293 cells transfected
45 with SV40 large antigen. COSA2 are a subclone of COS1 cells; subcloned in soft agar.

30 FuGENE Method: 293T cells were seeded at 2×10^5 cells per well of a 6 well culture
plate. Following overnight growth, cells were at approximately 40-50% confluency. Media
50 was changed a few hours before transfection (2 ml/well). For each sample, 3 μ l of FuGENE
6 Transfection Reagent (Roche Molecular Biochemicals, Indianapolis, IN) was diluted into

0.1 ml of serum-free culture medium (DME with 10 mM Hepes) and incubated at room temperature for 5 min. One microgram of DNA for each sample (0.5-2 mg/ml) was added to a separate tube. The diluted FuGENE reagent was added drop-wise to the concentrated DNA. After gentle tapping to mix, this mixture was incubated at room temperature for 15 minutes. The mixture was added dropwise onto the cells and swirled gently to mix. The cells were then incubated at 37 degrees C, in an atmosphere of 7.5% CO₂. The conditioned media and cells were harvested after 48 hours. Conditioned media was collected, centrifuged and isolated from the pellet. Protease inhibitors (5 µg/ml E64, 2 µg/ml pepstatin, 0.2 mM PMSF) were added prior to freezing. The cell monolayer was rinsed once with PBS, then 0.5 ml of lysis buffer (1 mM HEPES, pH 7.5, 1 mM EDTA, 0.5% Triton X-100, 1 mM PMSF, 10 µg/ml E64) was added. The lysate was frozen and thawed, vortex mixed, then centrifuged, and the supernatant was frozen until assayed.

Effective Method. DNA (0.6µg) was added with "EFFECTENE" reagent (Qiagen, Valencia, CA) into a 6-well culture plate using a standard transfection protocol according to manufacturer's instructions. Cells were harvested 3 days after transfection and the cell pellets were snap frozen. Whole cell lysates were prepared and various amounts of lysate were tested for β-secretase activity using the MBP-C125sw substrate. FIG. 14B shows the results of these experiments, in which picomoles of product formed is plotted against micrograms of COS cell lysate added to the reaction. The legend to the figure describes the enzyme source, where activity from cells transfected with DNA from pCEKclone27 and PCEKclone53 (clones 27 and 53) using Effective are shown as closed diamonds and solid squares, respectively, activity from cells transfected with DNA from clone 27 prepared with FuGENE are shown as open triangles, and mock transfected and control plots show no activity (closed triangles and "X" markers). Values greater than 700 pM product are out of the linear range of the assay.

Example 7

Preparation of P10-P4'sta(D->V) Sepharose Affinity Matrix

A. Preparation of P10-P4'sta(D->V) inhibitor peptide

P10-P4'sta(D->V) has the sequence NH₂-KTEEISEVN[sta]VAEF-COOH (SEQ ID NO: 72), where "sta" represents a statine moiety. The synthetic peptide was synthesized in a peptide synthesizer using boc-protected amino acids for chain assembly. All chemicals,

5 reagents, and boc amino acids were purchased from Applied Biosystems (ABI; Foster City, CA) with the exception of dichloromethane and N,N-dimethylformamide which were from Burdick and Jackson. The starting resin, boc-Phe-OCH₂-Pam resin was also purchased from
10 ABI. All amino acids were coupled following preactivation to the corresponding HOBt ester using 1.0 equivalent of 1-hydroxybenzotriazole (HOBt), and 1.0 equivalent of N,N-dicyclohexylcarbodiimide (DCC) in dimethylformamide. The boc protecting group on the
15 amino acid α -amine was removed with 50% trifluoroacetic acid in dichloromethane after each coupling step and prior to Hydrogen Fluoride cleavage.

Amino acid side chain protection was as follows: Glu(Bzl), Lys(Cl-CBZ),

10 Ser(OBzl), Thr(OBzl). All other amino acids were used with no further side chain protection including boc-Statine.

20 [(Bzl) benzyl, (CBZ) carbobenzoxy, (Cl-CBZ) chlorocarbobenzoxy, (OBzl) O-benzyl]

The side chain protected peptide resin was deprotected and cleaved from the resin by reacting with anhydrous hydrogen fluoride (HF) at 0°C for one hour. This generates the
25 fully deprotected crude peptide as a C-terminal carboxylic acid.

Following HF treatment, the peptide was extracted from the resin in acetic acid and lyophilized. The crude peptide was then purified using preparative reverse phase HPLC on a
30 Vydac C4, 330Å, 10µm column 2.2cm I.D. x 25cm in length. The solvent system used with this column was 0.1% TFA / H₂O ([A] buffer) and 0.1% TFA / CH₃CN ([B] buffer) as the mobile phase. Typically the peptide was loaded onto the column in 2% [B] at 8-10 mL/min.
20 and eluted using a linear gradient of 2% [B] to 60% [B] in 174 minutes.

35 The purified peptide was subjected to mass spectrometry, and analytical reverse phase HPLC to confirm its composition and purity.

40 25 B. Incorporation into Affinity Matrix

All manipulations were carried out at room temperature. 12.5 ml of 80% slurry of NHS-Sepharose (i.e. 10 ml packed volume; Pharmacia, Piscataway, NJ) was poured into a Bio-Rad EconoColumn (BioRad, Richmond, CA) and washed with 165 ml of ice-cold 1.0
45 mM HCl. When the bed was fully drained, the bottom of the column was closed off, and 5.0 ml of 7.0 mg/ml P10-P4'sta(D->V) peptide (dissolved in 0.1 M HEPES, pH 8.0) was added.
30 The column was capped and incubated with rotation for 24 hours. After incubation, the column was allowed to drain, then washed with 8 ml of 1.0 M ethanolamine, pH 8.2. An
50 additional 10 ml of the ethanolamine solution was added, and the column was again capped

5 and incubated overnight with rotation. The column bed was washed with 20 ml of 1.5 M sodium chloride, 0.5 M Tris, pH 7.5, followed by a series of buffers containing 0.1 mM EDTA, 0.2% Triton X-100, and the following components; 20 mM sodium acetate, pH 4.5
10 (100 ml); 20 mM sodium acetate, pH 4.5, 1.0 M sodium chloride (100 ml); 20 mM sodium borate, pH 9.5, 1.0 M sodium chloride (200 ml); 20 mM sodium borate, pH 9.5 (100 ml).
5 Finally, the column bed was washed with 15 ml of 2 mM Tris, 0.01% sodium azide (no Triton or EDTA), and stored in that buffer, at 4°C.
15

Example 8

10 Co-Transfection of Cells with β -secretase and APP

20 293T cells were co-transfected with equivalent amounts plasmids encoding APP^{sw} or wt and β -secretase or control β -galactoside (β -gal) cDNA using FuGene 6 Reagent, as described in Example 4, above. Either pCEKclone27 or pohCJ containing full length β -
25 15 secretase were used for expression of β -secretase. The plasmid construct pohCK751 used for the expression of APP in these transfections was derived as described in Dugan et al., JBC, 270(18) 10982-10989(1995) and shown schematically in FIG. 21. A β -gal control plasmid was added so that the total amount of plasmid transfected was the same for each condition.
30 β -gal expressing pCEK and pohCK vectors do not replicate in 293T or COS cells. Triplicate wells of cells were transfected with the plasmid, according to standard methods described above, then incubated for 48 hours, before collection of conditioned media and cells. Whole
35 cell lysates were prepared and tested for the β -secretase enzymatic activity. The amount of β -secretase activity expressed by transfected 293T cells was comparable to or higher than that expressed by CosA2 cells used in the single transfection studies. Western blot assays were
40 25 carried out on conditioned media and cell lysates, using the antibody 13G8, and A β ELISAs carried out on the conditioned media to analyze the various APP cleavage products.

While the invention has been described with reference to specific methods and
45 embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention. All patent and literature references referred to herein
30 are herein incorporated by reference.

Claims

5

10

15

20

25

30

35

40

45

50

55

5

IT IS CLAIMED:

10

1. A β -secretase enzyme protein purified to apparent homogeneity.

5

2. The purified β -secretase enzyme protein of claim 1, wherein the enzyme has been purified sufficiently so that its activity in cleaving the 695-amino acid isotype of β -amyloid precursor protein (β -APP) between amino acids 596 and 597 thereof is at least 10,000-fold greater than an activity exhibited by a solubilized but unenriched membrane fraction from human 293 cells.

15

10

3. The purified β -secretase enzyme protein of claim 1, characterized by a specific activity of at least about 0.2×10^5 nM/h/ μ g protein in an MBP-C125sw substrate assay.

20

4. The purified β -secretase enzyme protein of claim 3, wherein said specific activity is at least 1.0×10^5 nM/h/ μ g protein.

25

15

5. The purified β -secretase enzyme protein of claim 1, wherein said protein is fewer than 450 amino acids in length, comprising a polypeptide having the amino acid sequence SEQ ID NO: 70 [63-452].

30

20

6. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 70 [63-452].

35

7. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 69 [63-501].

40

25

8. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 67 [58-501].

45

30

9. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 68 [58-452].

50

10. The purified protein of any of claims 1-5, wherein said protein comprises a polypeptide having the amino acid sequence SEQ ID NO: 58 [46-452].

55

5

11. The purified protein of claim 10, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 74 [22-452].

10

12. The purified protein of claim 10, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 58 [46-452].

15

13. The purified protein of claim 10, wherein said protein is characterized by an N-terminus at position 46 with respect to SEQ ID NO: 2 and a C-terminus between positions 452 and 470 with respect to SEQ ID NO: 2.

20

14. The purified protein of claim 10, wherein said protein is characterized by an N-terminus at position 22 with respect to SEQ ID NO: 2 and a C-terminus between positions 452 and 470 with respect to SEQ ID NO: 2.

25

15

15. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 43 [46-501].

30

16. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 66 [22-501].

20

35

17. The purified protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 2 [1-501].

40

18. The purified protein of any of claims 1-5, wherein said protein has an N-terminal residue corresponding to a residue selected from the group consisting of residues 22, 46, 58 and 63 with respect to SEQ ID NO: 2 and a C-terminus selected from a residue between positions 452 and 501 with respect to SEQ ID NO: 2.

45

19. The purified protein of claim 18, wherein said C-terminus is between residue positions 452 and 470 with respect to SEQ ID NO: 2.

50

55

5

20. The purified protein of claim 1, wherein said protein is isolated from a mouse.

10

21. The protein of claim 20, wherein said polypeptide has the sequence SEQ ID NO: 65.

5 22. The purified protein of any of claims 1-21, wherein said protein is produced by a heterologous cell.

15

23. A crystalline protein composition formed from a purified β -secretase protein.

10

20

24. The crystalline protein composition of claim 23, wherein said purified protein is characterized by a binding affinity for the β -secretase inhibitor substrate P10-P4'sta D \rightarrow V which is at least 1/100 of an affinity exhibited by a protein having the amino acid sequence SEQ ID NO: 43 [46-501], when said proteins are tested for binding to said substrate under the

25

15 same conditions.

30

25. The crystalline protein composition of either of claims 23-24, wherein said composition is formed from a protein having a sequence selected from the group consisting of SEQ ID NO: 66 [22-501], SEQ ID NO: 43[46-501], SEQ ID NO: 74 [22-452], SEQ ID NO: 43 [46-452], and SEQ ID NO: 71 [46-419].

20

35

26. The crystalline protein composition of either of claims 23-24, wherein said composition is formed from a protein having a sequence selected from the group consisting of SEQ ID NO: 2 [1-501], SEQ ID NO: 59[1-452], and SEQ ID NO: 60 [1-420].

40

25

27. The crystalline protein composition of either of claims 23-24, wherein said composition is formed from a protein having an N-terminal residue corresponding to a residue selected from the group consisting of residues 22, 46, 58 and 63 with respect to SEQ ID NO: 2 and a C-terminus selected from a residue between positions 452 and 501 with respect to SEQ ID

45

30 NO: 2.

50

28. The crystalline protein of claim 27, wherein said C-terminus is between residue positions 452 and 470 with respect to SEQ ID NO: 2.

55

5

29. The crystalline protein composition of any of claims 23-28, wherein said protein is glycosylated.

10

30. The crystalline protein composition of any of claims 23-28, wherein said protein is deglycosylated.

15

31. The crystalline protein composition of any of claims 23-30, wherein said composition further includes a β -secretase substrate or inhibitor molecule.

10

20

32. The crystalline protein composition of claim 31, wherein said β -secretase inhibitor is a peptide having fewer than about 15 amino acids and comprises the sequence SEQ ID NO: 78 (VMXVAEF; P3-P4'X D->V), including conservative substitutions thereof.

25

33. The crystalline protein composition of claim 31, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 72 [P10-P4'sta D->V], including conservative substitutions thereof.

30

34. The crystalline protein composition of any claims 31-35, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 81 [EVMXVAEF], wherein X is hydroxyethylene or statine.

35

35. The crystalline protein composition of claim 31, wherein said β -secretase inhibitor is characterized by a K_i of no more than about 0.5 mM.

40

25

36. The crystalline protein composition of claim 31, wherein said β -secretase inhibitor is characterized by a K_i of no more than about 50 μ M.

45

37. An isolated protein, comprising a polypeptide that (i) is fewer than about 450 amino acid residues in length, (ii) includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, and (iii) exhibits β -secretase

50

55

5 activity, as evidenced by an ability to cleave a substrate selected from the group consisting of
the 695 amino acid isotype of beta amyloid precursor protein (BAPP) between amino acids
596 and 597 thereof, MBP-C125wt and MBP-C125sw.

10

5 38. The protein of claim 37, wherein said polypeptide includes the amino acid sequence of
SEQ ID NO: 75 [63-423].

15

39. The protein of claim 37, wherein said polypeptide has the sequence SEQ ID NO: 75 [63-
423].

10

20

40. The protein of claim 37, wherein said amino acid sequence is at least 95% identical to
SEQ ID NO: 58 [46-452].

25

15 41. The protein of claim 40, wherein said polypeptide has the sequence SEQ ID NO: 58 [46-
452].

30

42. The protein of claim 37, wherein said protein consists of a polypeptide having the
sequence SEQ ID NO: 58 [46-452].

35

20 43. The protein of claim 37, wherein said protein consists of a polypeptide having the
sequence SEQ ID NO: 74 [22-452].

40

25

44. The protein of any claims 37-43, wherein said protein is expressed by a heterologous
cell.

45

45. A composition comprising the protein of any claims 37-44 and a β -secretase substrate or
inhibitor molecule.

50

30 46. The composition of claim 45, wherein said β -secretase substrate is selected from the
group consisting of MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable
fragments thereof.

55

5 47. The composition of claim 46, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4'wt), SEVNLDAEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFDAEF, SEVKMAAEF, SEVNLAEEF, SEVKLAAEF;
10 SEVKMLAEF, SEVNLLAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.

15 48. The composition of claim 45, wherein said β -secretase inhibitor is a peptide having fewer than about 15 amino acids and comprises the sequence SEQ ID NO: 78 (VM[X]VAEF, where X is hydroxyethylene or statine), including conservative substitutions thereof.

10 49. The composition of claim 48, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 81 (VM[X]VAEF, where X is hydroxyethylene or statine).

20 50. The composition of claim 45, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 72 (P10-P4'sta D->V), including conservative substitutions thereof.

25 51. The composition of any claims 45 and 48-50, wherein said β -secretase inhibitor has a K_i of no more than about 1 μ M.

30 52. The composition of any claims 45 and 48-50, wherein said β -secretase inhibitor is labeled with a detectable reporter molecule.

35 53. An isolated mouse β -secretase protein enzyme having the sequence SEQ ID NO: 65.

40 54. An antibody which binds specifically to a purified β -secretase protein comprising a polypeptide that includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, wherein said antibody further
45 lacks significant immunoreactivity with a protein a sequence selected from the group consisting of SEQ ID NO: 2 [1-501] and SEQ ID NO: 43 [46-501].
50

- 5 55. The antibody of claim 54, wherein said antibody is reactive with a protein selected from the group consisting of SEQ ID NO: 66 [22-501], SEQ ID NO: 67 [58-501], SEQ ID NO: 69 [63-501], SEQ ID NO: 59 [1-452], SEQ ID NO: 74 [22-452], SEQ ID NO: 58 [46-452], SEQ ID NO: 68 [58-452] and SEQ ID NO: 70 [63-452].
- 10 5
- 15 56. An isolated nucleic acid, comprising a sequence of nucleotides that encodes a β -secretase protein that is at least 95% identical to a protein selected from the group consisting of SEQ ID NO: 66 [22-501], SEQ ID NO: 43[46-501], SEQ ID NO: 57 [1-419], SEQ ID NO: 74 [22-452], SEQ ID NO: 58 46-452], SEQ ID NO: 59 [1-452], SEQ ID NO: 60 [1-420], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 70 [63-452], SEQ ID NO: 75 [63-423], and SEQ ID NO: 71 [46-419], or a complementary sequence of any of such nucleotides, and specifically excluding a nucleic acid encoding a protein having the sequence SEQ ID NO: 2 [1-501].
- 20 25 15
- 30 57. The isolated nucleic acid of claim 56, wherein said sequence of nucleotides encodes a protease having an amino acid sequence SEQ ID NO: 58 [46-452].
- 35 58. The isolated nucleic acid of claim 56, wherein said sequence of nucleotides encodes a protease having the sequence SEQ ID NO: 43 [46-501].
- 40 20
- 45 59. The isolated nucleic acid of claim 56, wherein said sequence of nucleotides encodes a protease having the sequence SEQ ID NO: 66 [22-501].
- 50 25 60. The isolated nucleic acid of claim 56, wherein said sequence of nucleotides encodes a protease having the sequence SEQ ID NO: 74 [22-452].
- 55 45 61. A expression vector, comprising
30 the isolated nucleic acid of any of claims 56-60, and
operably linked to said nucleic acid, regulatory sequences effective for expression of
the nucleic acid in a selected host cell.

5

62. The recombinant expression vector of claim 61, wherein said vector is suitable for transfection of a bacterial cell.

10

5

63. A heterologous cell transfected with the vector of any of claims 61-62, wherein said cell expresses a biologically active β -secretase.

15

64. The cell of claim 63, wherein said cell is a eukaryotic cell.

10

65. The cell of claim 63, wherein said cell is a bacterial cell.

20

66. The cell of claim 63, wherein said cell is an insect cell.

25

15 67. The cell of claim 63, wherein said cell is a yeast cell.

68. A method of producing a recombinant β -secretase enzyme, comprising culturing a cell according to any of claims 63-67 under conditions to promote growth of said cell, and subjecting an extract or cultured medium from said cell to an affinity matrix.

30

20

69. The method of claim 68, wherein said affinity matrix contains a β -secretase inhibitor molecule.

35

70. The method of claim 69, wherein said inhibitor molecule is SEQ ID NO: 72 [P10-P4'staD->V].

40

25

71. The method of claim 68, wherein said matrix contains an antibody characterized by an ability to bind β -secretase.

45

30

72. The method of claim 71, wherein said antibody is according to any of claims 54-55.

50

55

5

73. A heterologous cell, comprising

10

(i) a nucleic acid molecule encoding an active β -secretase protein according to any of claims 37-43;

5

(ii) a nucleic acid molecule encoding a β -secretase substrate molecule; and

(iii) operatively linked to (i) and (ii), a regulatory sequence effective for expression of said nucleic acid molecules in said cell.

15

74. The cell of claim 73, wherein said nucleic acid encoding said β -secretase protein is heterologous to said cell.

20

75. The cell of claim 73, wherein both said nucleic acids encoding said β -secretase protein encoding said β -secretase substrate molecule are heterologous to said cell.

25

76. The cell of claim 73, wherein said β -secretase substrate molecule is selected from the group consisting of MBP-C125wt, MBP-C125sw, APPwt, APPsw, and β -secretase cleavable fragments thereof.

30

77. The cell of claim 76, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4'wt), SEVNLD AEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFDAEF, SEVKMAAEF, SEVNLA AEF, SEVKLAAEF; SEVKMLAEF, SEVNLLAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.

35

40

25

78. A method of screening for compounds that inhibit $A\beta$ production, comprising contacting an isolated β -secretase polypeptide according to claim 37 with (i) a test compound and (ii) a β -secretase substrate, and selecting the test compound as capable of inhibiting $A\beta$ production if said β -secretase polypeptide exhibits less β -secretase activity in the presence of said compound than in the absence of said compound.

45

30

50

55

5 79. The method of claim 78, wherein said active β -secretase polypeptide has a sequence selected from the group consisting of SEQ ID NO: 43 [46-501] and SEQ ID NO: 58 [46-452].

10 80. The method of claim 78, wherein said β -secretase polypeptide and said substrate are
5 produced by a cell according to claim 73.

15 81. The method of claim 78, which further includes administering said test compound to a mammalian subject having Alzheimer's disease or Alzheimer's disease-like pathology, and selecting said compound as a therapeutic agent candidate if, following such administration,
20 10 said subject maintains or improves cognitive ability or said subject shows reduced plaque burden.

25 82. The method of claim 81, wherein said subject is a mammalian species comprising a transgene.

15 30 83. The method of claim 81, wherein said subject is a mouse bearing a transgene which encodes a human β -amyloid precursor protein (β -APP), including a mutant variant thereof.

35 20 84. The method of any of claims 78-83, wherein said β -secretase substrate is selected from the group consisting of MBP-C125wt, MBP-C125sw, APP, APP_{sw}, and β -secretase-cleavable fragments thereof.

40 25 85. The method of claim 78, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4'wt), SEVNLDAEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFD AEF, SEVKMAAEF, SEVNLA AEF, SEVKLAAEF;
45 SEVKMLAEF, SEVNLLAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.

- 5 86. A method of screening for compounds that inhibit A β production, comprising measuring
binding of a purified β -secretase polypeptide according to any of claims 1-22 or 37-43 with
a β -secretase inhibitor compound in the presence of a test compound, and selecting the test
10 compound as β -secretase active-site binding compound, if binding of the inhibitor in the
5 presence of said test compound is less than binding of the inhibitor in the absence of said test
compound.
- 15 87. The method of claim 86, wherein said inhibitor compound is labeled with a detectable
marker.
- 10 88. The method of claim 86, wherein said β -secretase inhibitor is a peptide having fewer than
20 about 15 amino acids and comprises the sequence SEQ ID NO: 78 (VM[X]VAEF, where X
is hydroxyethylene or statine), including conservative substitutions thereof.
- 25 89. The method of claim 86, wherein said β -secretase inhibitor has the sequence SEQ ID
NO: 72 (P10-P4'sta D->V), including conservative substitutions thereof.
- 30 90. The method of claim 86, wherein said β -secretase inhibitor has a K_i with respect to β -
secretase of less than about 50 μ M.
- 20 91. A β -secretase inhibitor compound selected according to the method of any of claims 78-
35 80.
- 40 92. The inhibitor of claim 91, wherein said compound is selected from a phage display
25 selection system.
- 45 93. The compound of claim 92, wherein the phage display selection system is biased for the
sequence SEQ ID NO: 72 [P10-P4'D->V].
- 50

5 94. A β -secretase inhibitor compound selected according to the method of any of claims 81-90.

10 95. The inhibitor of claim 94, wherein said compound is selected from a phage display
5 selection system.

15 96. The compound of claim 95, wherein the phage display selection system is biased for the sequence SEQ ID NO: 72 [P10-P4'D \rightarrow V].

20 97. A β -secretase inhibitor compound selected according to the method of any of claims 86-90.

25 98. The inhibitor of claim 97, wherein said compound is selected from a phage display selection system.

30 99. The compound of claim 98, wherein the phage display selection system is biased for the sequence SEQ ID NO: 72 [P10-P4'D \rightarrow V].

35 100. A β -secretase inhibitor, comprising a peptide containing the sequence SEQ ID NO: 78
20 (VM[X]VAEF, where X is hydroxyethylene or statine), including conservative substitutions thereof.

40 101. The β -secretase inhibitor of claim 100, having the sequence SEQ ID NO: 72 (P10-P4'sta D \rightarrow V).

45 25 102. The β -secretase inhibitor of claim 100, having the sequence SEQ ID NO: 78.

50 103. The β -secretase inhibitor of claim 100, having the sequence SEQ ID NO: 81.

5

104. A screening kit, comprising

10

an isolated β -secretase protein according to any of claims 1- 22 or 37-43,

a cleavable β -secretase substrate, and

5

means for detecting cleavage of said substrate by β -secretase.

15

105. The screening kit of claim 104, wherein said β -secretase protein is present in a heterologous cell.

20

106. The screening kit of claim 104, wherein said β -secretase substrate molecule is selected from the group consisting of MBP-C125wt, MBP-C125sw, APPwt, APPsw, and β -secretase cleavable fragments thereof.

25

107. The screening kit of claim 106, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4'wt), SEVNLDAEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFDAEF, SEVKMAAEF, SEVNLAAEF, SEVKLAAEF; SEVKMLAEF, SEVNLLAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.

35

20

108. A knock-out mouse, characterized by inactivation or deletion of an endogenous β -secretase gene.

40

109. The knock-out mouse of claim 108, wherein said β -secretase gene encodes a protein having at least 90% sequence identity to the sequence SEQ ID NO: 65.

45

110. The knock-out mouse of claim 108, wherein said deletion is inducible.

50

55

5

111. The knock-out mouse of claim 110, wherein said inducible expression is effected by a Cre-lox expression system inserted into the mouse genome.

10

5 112. A method of screening for drugs effective in the treatment of Alzheimer's disease or other cerebrovascular amyloidosis characterized by A β deposition, comprising

15

administering to a mammalian subject characterized by overexpression of β -APP and/or deposition of A β a test compound selected for its ability to inhibit β -secretase activity a β -secretase protein according to any of claims 37-43, and

20

10 selecting the compound as a potential therapeutic drug compound, if it reduces the amount of A β deposition in said subject or if it maintains or improves cognitive ability in said subject.

25

113. The method of claim 112, wherein said mammalian subject is a transgenic mouse

15 bearing a transgene encoding a human β -APP or a mutant thereof.

30

114. A method of treating a patient afflicted with or having a predilection for Alzheimer's disease or other cerebrovascular amyloidosis, comprising

35

20 blocking the enzymatic hydrolysis of APP to A β in the patient by administering to the patient a pharmaceutically effective dose of a compound effective to inhibit a β -secretase enzyme protein according any of claims 37-43.

40

115. The method of claim 114, wherein said compound is derived from a peptide selected

25 from the group consisting of SEQ ID NO: 72, SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 97.

45

50

55

5

116. A method of inhibiting enzymatic proteolysis of APP to A β in a tissue, comprising contacting said tissue with a compound effective to inhibit the enzymatic activity of a β -secretase protein according to any of claims 37-43.

10

117. The method of claim 116, wherein said inhibition of enzymatic activity is evidenced by a K_i of less than about 50 μ M in a MBP-C125sw assay.

15

118. A therapeutic drug composition for the treatment of Alzheimer's disease or other cerebrovascular amyloidosis characterized by deposition of A β peptide, wherein the active compound in said drug is selected for its ability to inhibit the enzymatic activity of a β -secretase protein according to claim 37.

20

25

119. The therapeutic drug of claim 118, wherein said inhibition of enzymatic activity is evidenced by a K_i of less than about 50 μ M in a MBP-C125sw assay.

15

30

120. The therapeutic drug of claim 118, wherein said drug is derived from a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 97.

20

35

121. A method of diagnosing the presence of or a predilection for Alzheimer's disease in a patient, comprising

40

detecting the expression level of a gene comprising a nucleic acid encoding β -secretase in a cell sample from said patient, and
diagnosing the patient as having or having a predilection for Alzheimer's disease, if said expression level is significantly greater than a pre-determined control expression level.

25

45

122. The method of claim 121, wherein said gene comprises a nucleic acid that encodes a β -secretase protein that is at least 95% identical to a protein selected from the group consisting of SEQ ID NO: 66 [22-501], SEQ ID NO: 43[46-501], SEQ ID NO: 57 [1-419], SEQ ID NO: 74 [22-452], SEQ ID NO: 58 46-452], SEQ ID NO: 59 [1-452], SEQ ID NO: 60 [1-

30

50

55

- 5 420], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 69 [63-501], SEQ ID
NO: 70 [63-452], SEQ ID NO: 75 [63-423], and SEQ ID NO: 71 [46-419], or a
complementary sequence of any of such nucleotides.
- 10 5 123. The method of claim 121, wherein said nucleic acid specifically excludes a nucleic acid
encoding a protein having the sequence SEQ ID NO: 2 [1-501].
- 15 124. The method of claim 121, wherein said nucleic acid encodes a protein having the
sequence SEQ ID NO: 2 [1-501].
- 10 125. The method of claim 121, wherein said measuring is carried out in a whole cell assay.
- 20 126. The method of claim 109, wherein said measuring is carried out on a nucleic acid
derived from a cell sample of said patient.
- 25 15 127. A method of purifying a β -secretase protein enzyme molecule, comprising
contacting an impure sample containing β -secretase enzyme activity with an affinity
matrix which includes a β -secretase inhibitor.
- 30 20 128. The method of claim 127, wherein said β -secretase inhibitor comprises a peptide having
the sequence SEQ ID NO: 78 (VM[X]VAEF, where X is hydroxyethylene or statine),
including conservative substitutions thereof.
- 35 129. The method of claim 128, wherein said β -secretase inhibitor has the sequence SEQ ID
25 NO: 72 (P10-P4'sta D->V).
- 40 130. The method of claim 128, wherein said β -secretase inhibitor has the sequence SEQ ID
NO: 78.
- 45 30 131. The method of claim 128, wherein said β -secretase inhibitor has the sequence SEQ ID
NO: 81.

ATGGCCCAAGCCCTGCCCTGGCTCCTGCTGTGGATGGGCGCGGGAGT
GCTGCCTGCCACGGCACCCAGCACGGCATCCGGCTGCCCCTGCGCA
GCGGCCTGGGGGGCGCCCCCTGGGGCTGCGGCTGCCCCGGGAGAC
CGACGAAGAGCCCGAGGAGCCCGGCCGGAGGGGCAGCTTTGTGGAGA
TGGTGGACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAG
ATGACCGTGGGCAGCCCCCGCAGACGCTCAACATCCTGGTGGATACA
GGCAGCAGTAACCTTTCAGTGGGTGCTGCCCCCACCCTTCCTGCAT
CGCTACTACCAGAGGCAGCTGTCCAGCACATACCGGGACCTCCGGAAG
GGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAAGGGGAGCTGGG
CACCGACCTGGTAAGCATCCCCATGGCCCCAACGTCACTGTGCGTGC
CAACATTGCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCC
AACTGGGAAGGCATCCTGGGGCTGGCCTATGCTGAGATTGCCAGGCCT
GACGACTCCCTGGAGCCTTTCTTTGACTCTCTGGTAAAGCAGACCCACG
TTCCCAACCTCTTCTCCCTGCAGCTTTGTGGTGCTGGCTTCCCCCTCAA
CCAGTCTGAAGTGCTGGCCTCTGTCCGAGGGAGCATGATCATTGGAGG
TATCGACCACTCGCTGTACACAGGCAGTCTCTGGTATACACCCATCCGG
CGGGAGTGGTATTATGAGGTGATCATTGTGCGGGTGGAGATCAATGGA
CAGGATCTGAAAATGGACTGCAAGGAGTACAACATGACAAGAGCATTG
TGGACAGTGGCACCACCAACCTTCGTTTGCCCAAGAAAGTGTTTGAAGC
TGCAGTCAAATCCATCAAGGCAGCCTCCTCCACGGAGAAGTTCCTGAT
GGTTTCTGGCTAGGAGAGCAGCTGGTGTGCTGGCAAGCAGGCACCACC
CCTTGGAACATTTTCCAGTCATCTCACTCTACCTAATGGGTGAGGTAC
CAACCAAGTCCCTCCGCATCACCATCCTTCCGCAGCAATACCTGCGGCCA
GTGGAAGATGTGGCCACGTCCCAAGACGACTGTTACAAGTTTGCCATCT
CACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTT
CTACGTTGTCTTTGATCGGGCCCGAAAACGAATTGGCTTTGCTGTCAGC
GCTTGCCATGTGCACGATGAGTTCAGGACGGCAGCGGTGGAAGGCCCT
TTTGTACCTTGGACATGGAAGACTGTGGCTACAACATTCCACAGACAG
ATGAGTCAACCCTCATGACCATAGCCTATGTCATGGCTGCCATCTGCGC
CCTCTTCATGCTGCCACTCTGCCTCATGGTGTGTCAGTGGCGCTGCCTC
CGCTGCCTGCGCCAGCAGCATGATGACTTTGCTGATGACATCTCCCTGC
TGAAG

FIG. 1A

2/26

CCATGCCGGCCCCCTCACAGCCCCGCCGGGAGCCCCGAGCCCCGCTGCCAGGCTGGC
CGCCGCSGTGCCGATGTAGCGGGCTCCGGATCCCAGCCTCTCCCCCTGCTCCCGTGC
TCTGCGGATCTCCCCTGACCGCTCTCCACAGCCCCGACCCGGGGGCTGGCCCCAGG
GCCCTGCAGGCCCTGGCGTCCTGATGCCCCAAGCTCCCTCTCCTGAGAAGCCACC
AGCACCACCCAGACTTGGGGGCAGGCGCCAGGGACGGACGTGGGGCCAGTGCGAGC
CCAGAGGGCCCCGAAGGCCGGGGCCCCACCATGGCCCCAAGCCCTGCCCTGGCTCCTG
CTGTGGATGGGCGCGGGAGTGCTGCCTGCCACGGCACCCAGCACGGCATCCGGC
TGCCCCCTGCGCAGCGGCCCTGGGGGGCGCCCCCTGGGGCTGCGGCTGCCCGGG
AGACCGACGAAGAGCCCCGAGGAGCCCCGGCCGGAGGGGCAGCTTTGTGGAGATGGT
GGACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACCGTGGGC
AGCCCCCGCAGACGCTCAACATCCTGGTGGATACAGGCAGCAGTAACCTTTCAGT
GGGTGCTGCCCCCAACCCCTTCTGTCATCGCTACTACCAGAGGCAGCTGTCCAGCA
CATACCGGGACCTCCGGAAGGGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAA
GGGGAGCTGGGCACCGACCTGGTAAGCATCCCCCATGGCCCCAACGTCACCTGTGCG
TGCCAACATTGCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCCAACCTG
GAAGGCATCCTGGGGCTGGCCTATGCTGAGATTGCCAGGCCTGACGACTCCCTGGA
GCCTTTCTTTGACTCTCTGGTAAAGCAGACCCACGTTCCCAACCTCTTCTCCCTGCAG
CTTTGTGGTGCTGGCTTCCCCCTCAACAGTCTGAAGTGCTGGCCTCTGTGCGGAGG
GAGCATGATCATTGGAGGTATCGACCACTCGCTGTACACAGGCAGTCTCTGGTATAC
ACCCATCCGGCGGGAGTGATTATGAGGTGATCATTGTGCGGGTGGAGATCAATG
GACAGGATCTGAAAATGGACTGCAAGGAGTACAACCTATGACAAGAGCATTGTGGACA
GTGGCACCACCAACCTTCGTTTGCCCAAGAAAGTGTTTGAAGCTGCAGTCAAATCCA
TCAAGGCAGCCTCCTCCACGGAGAAGTTCCTGATGGTTTCTGGCTAGGAGAGCAG
CTGGTGCTGGCAAGCAGGCACCAACCCCTTGGAAACATTTCCAGTCATCTCACTC
TACCTAATGGGTGAGGTTACCAACCACTCCTTCCGCATCACCATCCTTCCGCAGCAA
TACCTGCGGCCAGTGGAAGATGTGGCCACGTCCCAAGACGACTGTTACAAGTTTGCC
ATCTCACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTTCTAC
GTTGTCTTTGATCGGGCCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGCCATGTG
CACGATGAGTTCAGGACGGCAGCGGTGGAAGGCCCTTTTGTACCTTGGACATGGA
AGACTGTGGCTACAACATTCCACAGACAGATGAGTCAACCCCTCATGACCATAGCCTA
TGTGATGGCTGCCATCTGCGCCCTCTTCATGCTGCTGCACTCTGCCTCATGGTGTGTCA
GTGGCGCTGCCTCCGCTGCCTGCGCCAGCAGCATGATGACTTTGCTGATGACATCT
CCCTGCTGAAGTGAGGAGGCCCATGGGCAGAAGATAGAGATTCCCCTGGACCACAC
CTCCGTGGTTCACTTTGGTCACAAGTAGGAGACACAGATGGCACCTGTGGCCAGAG
CACCTCAGGACCCCTCCCCACCCACCAATGCCTCTGCCTTGATGGAGAAGGAAAAG
GCTGGCAAGGTGGGTTCCAGGGACTGTACCTGTAGGAAACAGAAAAGAGAAGAAAG
AAGCACTCTGCTGGCGGGAATACTCTTGGTCACCTCAAATTTAAGTCGGGAAATTCT
GCTGCTTGAAACTTCAGCCCTGAACCTTTGTCCACCATTCTTTAAATTCTCCAACCC
AAAGTATTCTTCTTTCTAGTTTCAGAAGTACTGGCATCACACGCAGGTTACCTTGG
CGTGTGTCCCTGTGGTACCCTGGCAGAGAAGAGACCAAGCTTGTTTCCCTGCTGGC
CAAAGTCAGTAGGAGAGGATGCACAGTTTGCTATTTGCTTTAGAGACAGGGACTGTA
TAAACAAGCCTAACATTGGTGCAAGATTGCCTCTTGAATT

FIG. 1B

MAQALPWLLLWMGAGVLP AHGTQH GIRLPLRSGLG G APLGLRLP
RETDEEPEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPQT
LNILVDTGSSNF AVGAAPHPFLH RYYQRQLSSTYRDLRKGVVYPY
TQGWEGELGTDLV SIPHGPNVTVRANIAAITESDKFFINGSNWE
GILGLAYAEIARPDDSLEPFFDSL VKQTHVPNLFSLQLCGAGFPLN
QSEVLASVGGSMIIGGIDHSLYTGSLWYTPIRREWYYEVIIVRVEIN
GQDLKMDCKEYNYDKSIVDSGTTNLRLPKKVFEAAVKSIIKAASST
EKFPDGFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRT
ILPQQYLRPVEDVATSQDDCYKFAISQSSTGTVMGAVIMEGFYVV
FDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQ
TDESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRQQHDDF
ADDISLLK

FIG. 2A

4/26

ETDEEPEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSPQT
LNILVDTGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVYPY
TQGWEGELGTDLVSIHPGNVTVRANIAAITESDKFFINGSNWE
GILGLAYAEIARPDDSLEPFFDSL VKQTHVPNLFSLQLCGAGFPLN
QSEVLASVGGSMIIGGIDHSLYTGSLWYTPIRREWYYEVIIVRVEIN
GQDLKMDCKEYNYDKSIVDSGTTNLRPKKVFEAAVKSIIKAASST
EKFPDGFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRIT
ILPQQYL RPVEDVATSQDDCYKFAISQSSTGTVMGAVIMEGFYV
FDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQ
TDESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRQQHDDF
ADDISLLK

FIG. 2B

5/26


FIG. 3A

MAQALPWLLWLGAGVLPAGHTQHGIPLRSLGGLRLPRETDEEPE
EPGRRGSFVEMVDNLRGKSGQGYVEMT
VGSPQTNLNLDVTGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVPYPT
QGWEGELGTDLVSIHPGPNVTVRANI
AAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFFDSL VKQTHV PNL FSLQL
CGAGFPLNQSEVLASVGGSMIIGGI
DHSLYTGSLWYTPIRREWYVEIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNL
RLPKKVFEAAVKSIIKASSTEKFPD
GFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRTILPQQYLRPVEDVA
TSQDDCYKFAISQSSTGTVMGAVIME
GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPVTLDMEDCGYNIPQTDED
YKDDDDK

FIG. 3B

ETDEEPEEPGRRGSFVEMVDNLRGKSGQGYVEMT
VGSPQTNLNLDVTGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVPYPT
QGWEGELGTDLVSIHPGPNVTVRANI
AAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFFDSL VKQTHV PNL FSLQL
CGAGFPLNQSEVLASVGGSMIIGGI
DHSLYTGSLWYTPIRREWYVEIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNL
RLPKKVFEAAVKSIIKASSTEKFPD
GFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRTILPQQYLRPVEDVA
TSQDDCYKFAISQSSTGTVMGAVIME
GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPVTLDMEDCGYNIPQTDED
YKDDDDK

6/26

**FIG. 4**




FIG. 5

8/26




FIG. 6A

FIG. 6B

9/26




FIG. 7

SP flow thru
SP load
load
Flow thru
Eluate 1
Eluate 2
Eluate 3
293T standard



FIG. 8

10/26




FIG. 9

11/26




FIG. 10

12/26

CTGTTGGGCTCGCGGTTGAGGACAAACTCTTCGCGGTCTTTCCAGTACTCT
 TGGATCGGAAACCCGTGGCCTCCGAACGGTACTCCGCCACCGAGGGACCT
 GAGCGAGTCCGCATCGACCGGATCGGAAAACCTCTCGACTGTTGGGGTGAG
 TACTCCCTCTCAAAAGCGGGCATGACTTCTGCGCTAAGATTGTCAGTTTCC
 AAAAACGAGGAGGATTTGATATTCACCTGGCCCCGCGGTGATGCCTTTGAGG
 GTGGCCGCGTCCATCTGGTCAGAAAAGACAATCTTTTTGTGTCAAGCTTG
 AGGTGTGGCAGGCTTGAGATCTGGCCATACACTTGAGTGACAATGACATCC
 ACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTCCAAGTGCAGGTG
 ACTCTAGACCC

FIG. 11A




FIG. 11B

13/26




FIG. 12

FIG. 13A

1 TTCTCATGTTTGACAGCTTATCTATGCCAGATCCCGCCAAAGTTCCTGCATTCCTCCACAGCCAGAACTGGTAGGTAATGAGATCCGATGTACGGCCAGATATAC

107 GGCTGCACATTGATTTACTGACTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATACGCCATATATGAGATTCGGCGTACATAAATCTACGGTAAATGGCC

213 CCGGCTGGCTGACCGCCCAACGACCCCGGCCCATTTGAAGTCAATATGACGATATGTCCTCATATGAAACCCCATAGGACTTTCCATTTGAGCTCAATGGGTGGAGCT

319 ATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGATTCATATGCCAAGTAGGCCCGCATTTGAAGTCAATGACGGTAAATGGCGGCCCTGGCAATATGCCCA

425 GTACATGACCTTATGGGACTTTCTACTTGGCAGTACATCAAGTATTAGTCATGCTATTAACCATGGTATGGCGTTTGGCGATACATCAATGGCGTGGATAG

531 CCGTTTGCATCAGCGGATTTCCCAAGTCTCCACCCATTTGACGGTCAATGGGAGTTGTGTTTGGCACCAAAATCAACGGGACTTTCCCAAAATGTGTAACTACCTCG

637 CCCCATTGACGCAATATGGCGGTATGGCGTGTACGGTGGGAGGTCATATTAAGCAGAGCTCTTGGCTAACTAGAGAACCCATCTCTACTTGGCTTATGGAAATTAA

743 TAGACTCACTATAGGAGACCCCAAGCTCTCTTGGGCTGGCGGTGTAGGACAAACTCTTTCGGCGTCTTTCCAGTACTCTTGGATCGGAAACCGGCTGGGCTCCGAA

849 CCGTACTCCCGCACCGGAGGCTTGAGCGAGTCCGCATGACCCGATCCGAAAAAGCTCTGGAGCTTTGGGCTGAGTACTGCTCTCAAAAGCGGCCATGACTTCTG

955 CCGTAAGATTGTGAGTTTCCAAAAAGCAGGAGATTTCATATTCATCTGGCGCCGGGTGATGCTTTGAGGGTGGCCCGGCTGCATCTGGTCAGAAAGACAAATCTT

1061 TTTGTGTCAAGCTTGAGGTGTCCAGAGCTTGAGATCTGGCATACACTTGAGTGCAATGACATCCACTTTGCCCTTCTCTCCACAGGCTCCCACTCCACGGTCC

1167 AACTGCACTGCACTCTAGACCCGGGAAATCTGCAGATATCCATCACATGCCGCCAATCTGTGCCACGCCCGCCGGGAGCTGGAGACCCCGAGCTGCATTTATGG

1273 TGCGCTGAGCACCGAACCGACGCCAGGAGCCCGGAGCCCTTGCGCCAGGCTGGCCCGCGCTCCGATGTAGAGGGCTCCGATGCCAGCTCTGCCCTGCTCCCGTCTCTG

1379 CCGTCCACAGCCCGCGGAGCGCCGCCCGCTGCCAGGCTGGCCCGCGCTCCGATGTAGAGGGCTCCGATGCCAGCTCTGCCCTGCTCCCGTCTCTG

1485 CCGATCTCCCGTGACCGCTCTCCACAGCCCGGAGCCCGGGGCTGGCCAGGCGCTCCGAGCGCTGGCGTCTGATGCCCGCCCGAGCTCCCTCTCTGAGAAAGCCAC

1591 CAGCAACCCAGACTTTGGGGCGAGGGGCCAGGAGCGGAGCTGGGGCAGTGGCAGCCAGAGGCGCCGAGAGCCCGGGCCACCTATG GGC CAA GGC CTG
31Met Ala Gln Ala Leu

1690 CCG TGG CTC CTG CTG TGG ATG GCG GCG GSA GTG CTG CCT GCG CAC GCG ACC CAG CAC GGC ATC GCG CTG CCG CTG GCG
61Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg

1768 ACC GCG CTG GCG GCG GCG GCG CTG GCG CTG GCG CTG GCG ACC CAG GAA GAG CCG CAG GAG CCG CCG GCG AGG
321Ser Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Gly Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg

1846 GCG ACC TTT GTG CAG ATG GTG CAC MAC CTG AGG GCG MAG TCG GCG CAG GCG TAC TAC GTG GAG ATG ACC GTG GCG ACC
581Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser

1924 CCG CCG CAG AGC CTC AAC ATC CTG GTG GAT ACA GCG AGC AGT AAC TTT GCA GTG GGT GCT GCG CCC CAC CCC TTC CTG
841Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Ala Glu Pro His Pro Phe Leu

2002 CAT GCG TAC TAC CAG AGG CAG CTG TCC AGC ACA TAC CCG CAG CTC CCG AAG GGT GTG TAT GTG CCC TAC ACC CAG GCG
1101His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly

2080 AAG TCG GAA GCG CAG CTG GCG ACC CAG CTG GTA ACC ATC CCC CAT GCG CCC AAC GTC ACT GTG CGT GCG AAC ATT GCT
1361Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro Phe His Glu Pro Asn Val Thr Val Arg Ala Asn Ile Ala

2158 GCG ATC ACT GAA TCA CAG AAG TTC TTC ATC AAC GCG TCC AAC TGC GAA GGC ATC CTG GCG CTG GCG TAT GCT GAG ATT
1621Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile

2236 GCG AGG CCT GAC GAC TCC CTG GAG CTT TTC TTT GAG TCT CTC GTA AAG CAG ACC CAG GTT ACC AAC CTC TCC TCC CTG
1881Ala Arg Pro Asp Asp Thr Leu Glu Pro Phe Phe Asp Ser Thr Gln Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu

2314 CAG CTT TGT GGT GCT GCG TTC CCG CTC AAC CAG TCT GAA GTG CTG GCG TCT GTC GSA GCG ACC ATG ATC ATT GGA GGT
2141Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly

2392 ATC CAG CAC CTC CTG TAC ACA GCG AGT CTC TGG TAT ACA CCG ATC CCG GCG GAG TGG TAT TAT GAG CTC ATT ATT GTG
2401Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val

2470 GCG GTG CAG ATC AAT GCA CAG GAT CTG AAA ATG CAG TCG AAG GAG TAC AAC TAT GAG ACC AAT ATT GAG ACC GGT GCG
2661Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly

2548 ACC ACC AAC CTT GGT TGT CCG AAG AAA GTG TTT GAA GCT GCA GTC AAA TCC AAC GCA GCG TCC TCC ACC GAG AAG
2921Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys

2626 TTC CTT GAT GGT TTC TCG CTA GGA GAG CAG CTG GTG TGC TCA GCA GCG ACC ACC CTT TCG AAC ATT TTC CTA GCT
3181Phe Pro Asp Gly Phe Trp Leu Glu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Pro Trp Asn Ile Phe Cys Phe Val

15/26

FIG. 13B

[illegible]

[illegible]

17/26

FIG. 13D

[illegible]

18/26

FIG. 13E

15041 CTGCCAACTGTGATGGACACACCGTCAGTGGTCCGTCCGCCAGCCTCTCGATGAGCTGATCCTTGGGCGAGGACTGCCCGAGTCCGCCACCTGTGCAC
15147 GGGGATTTGGGCTCCACAAATCTCTGACGGACAAATGCCGCCATAACAGCGGTCATTGACTGCAAGGAGGGCATGTTGGGCGATTCCCAATAGGAGTCCGCCAACA
15253 TCTTCCTCTGGAGGCGGTGGTTGGGCGGTATGGAGCAGCAGACCGCCTACTTCCAGGGAGGCAATCCGAGCTTCCAGGATCCCGCGCGCTCCCGCGGTATATGCT
15359 CCGCATTTGCTCTTGACCAACTCTATCAGACCTTGGTTGACGGCAATTTCCATCATGCACTTGGGCCAGGCTCGATGGACCCAAATGGTCCCATCCGAGCGCGG
15465 ACTGTCCGGGTACACAAATCCCGCCAGAGCGCGCGCGCTCTCGACCGATGGCTGTGTAGAGTACTCCCGCATAGTGGAAACGGGAGATGGGGAGGCTAAGTG
15571 AATACCGGAAGGAGCAATACCGGAGGMAACCGCGCTATGACGGCAATAAAGACAGATAAAGCCAGCGGTGTGGGTGGTTTGTTCATAAAGCGCGGTTC
15677 GGTCCAGCGCTGGCACTCTGTGATACCCAGGAGACCCCATTTGGGCGAATAGCGCGCGGTTCCTTCCTTTTCCAGCCCAACCGCGCAAGTTCGGGTGAAGG
15783 CCGAGGCTCCAGCCAGGTGGGGCGCGAGCGCGCTGCCATAGCCACTGGCGCGGTGGGTAGGGAGCGGTCCCGCATGGGAATGGTTAATGTTGGTGGCGG
15889 TATATATTTGGGGTGGGTGGGTCTGGTCCAGACTGGACTGAGCAGACAGACCCATGGTTTGGATGGCTGGCATGGACGGCATGACTGGCGGACAC
15995 GATCAACCGGGGTCTGTGCTGCCAATACACCGGACCGCGCAAAACACCGCGGGATTCTGGGGTGGCAAGCTAGTGGACAA

Sall

→

19/26

CTGTTGGGCTCGCGGTTGAGGACAAACTCTTCGCGGTCTTTCCAGTACTCTTGGATCGGAAAC
 CCGTCGGCCTCCGAACGGTACTCCGCCACCGAGGGACCTGAGCGAGTCCGCATCGACCGGAT
 CGGAAAACCTCTCGACTGTTGGGGTGAGTACTCCCTCTCAAAAGCGGGCATGACTTCTGCGCT
 AAGATTGTCAAGTTTCCAAAAACGAGGAGGATTTGATATTACCTGGCCCGCGGTGATGCCTTT
 GAGGGTGGCCCGTCCATCTGGTCAGAAAAGACAATCTTTTGTGTCAAGCTTGAGGTGTGG
 CAGGCCTTGAGATCTGGCCATACATGAGTGACAATGACATCCACTTTGCCTTTCTCTCCACAG
 GTGTCCACTCCCAGGTCCAACCTGCAGGTCGACTCTAGACCC

FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B




FIG. 16A

FIG. 16B

22/26





FIG. 17A

FIG. 17B

23/26

**FIG. 18**

24/26




FIG. 19A

Wild-Type SequenceVal-Lys-Met-Asp...
Swedish SequenceVal-Asn-Leu-Asp...

FIG. 19B

25/26




FIG. 20

26/26




FIG. 21

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (11/15/2011)